

Heimildaskrá

Abramowitz, S.A., Flattery, D., Frances, K. og Berry, L. (2010). Linking a motivational interviewing curriculum to the chronic care model. *J Gen Intern Med*, 25(4), 620-626. doi: 10.1007/s11606-010-1426-6

Agnarsdóttir, A., Eyþórsdóttir, Á., Einarsson, E., Daniëlsdóttir, G., Kristjánsdóttir, H., Sverrisdóttir, K., ... Ólafsdóttir, H. (2011). *Punglyndi og kvíði–Klinískar leiðbeiningar*. Reykjavík: Landspítalinn.

AHCPR. (1993). *Depression Guideline Panel: Depression in primary care:2. Treatment of major depression: Clinical practice guideline*. Washington: U.S. Government Printing office.

Ahlström, B. H., Skärsäter, I., og Danielson, E. (2009). Living with major depression: Experiences from families' perspectives. *Scandinavian Journal of Caring Sciences*, 23(2), 309-316. doi:<http://dx.doi.org/10.1111/j.1471-6712.2008.00624.x>

Aina, Y., og Susman, J. L. (2006). Understanding comorbidity with depression and anxiety disorders. *The Journal of the American Osteopathic Association*, 106(2), 9-14.

Allen, J., og Annells, M. (2009). A literature review of the application of the geriatric depression scale, depression anxiety stress scales and post-traumatic stress disorder checklist to community nursing cohorts. *Journal of Clinical Nursing*, 18(7), 949-959. doi: <http://dx.doi.org/10.1111/j.1365-2702.2008.02731.x>

Andrews, G., (2008). Reducing the burden of depression. *Canadian Journal of Psychiatry*, 53(7), 420-7.

Anacker, C., Zunszain, P. A., Carvalho, L. A., og Pariante, C. M. (2011). The glucocorticoid receptor: Pivot of depression and of antidepressant treatment? *Psychoneuroendocrinology*, 36(3), 415-425. doi:<http://dx.doi.org/10.1016/j.psyneuen.2010.03.007>

Argyropoulos, S. V., og Wilson, S. J. (2005). Sleep disturbances in depression and the effects of antidepressants. *International Review of Psychiatry*, 17(4), 237-245.

Arna Skúladóttir, Auður Ragnarsdóttir og Elísabet Konráðsdóttir. (2005). Erum við föst í viðjum vanans? *Tímarit hjúkrunarfræðinga*, 81(4), 34-39.

Ay-woan, P., Sarah, C. P., Lyinn, C., Tsyr-jang, C., og Ping-chuan, H. (2006). Quality of life in depression: Predictive models. *Quality of Life Research*, 15(1), 39-48. doi:<http://dx.doi.org/10.1007/s11136-005-0381-x>

Bandura, A. (1982). Self-efficacy Mechanism in Human Agency. *American Psychologist*, 37(2), 122-147.

- Beautrais, A. L. (2003). Subsequent mortality in medically serious suicide attempts: A 5 year follow-up. *The Australian and New Zealand Journal of Psychiatry*, 37(5), 595-599.
- Beazley, P., og Gudjonsson, G. (2011). Motivating inpatients to engage with treatment: The role of depression and ward atmosphere. *Nordic Journal of Psychiatry*, 65(2), 95-100.
doi:<http://dx.doi.org/10.3109/08039488.2010.502244>
- Beck, A. T., MD. (2008). The evolution of the cognitive model of depression and its neurobiological correlates. *The American Journal of Psychiatry*, 165(8), 969-77.
- Bellino, S., Zizza, M., Rinaldi, C., og Bogetto, F. (2006). Combined treatment of major depression in patients with borderline personality disorder: A comparison with pharmacotherapy. *Canadian Journal of Psychiatry*, 51(7), 453-60.
- Belmaker, R. H., og Agam, G. (2008). Major depressive disorder. *The New England Journal of Medicine*, 358(1), 55-68.
- Bewernick, B. H., Hurlemann, R., Matusch, A., Kayser, S., Grubert, C., Hadrysiewicz, B., . . . Schlaepfer, T. E. (2010). Nucleus accumbens deep brain stimulation decreases ratings of depression and anxiety in treatment-resistant depression. *Biological Psychiatry*, 67(2), 110-116.
- Björgvin Ingimarsson. (2010). *Próffræðilegt mat á DASS sjálfsmatskvarðanum þunglyndi, kvíði og streita*. Sótt af
<http://skemman.is/item/view/1946/5411;jsessionid=48286F86D88F22A7A65FAD12DE54484C>
- Bjørnsgaard, J.H. (2007). SAMDATA Sektorrapport for det psykiske helsevernet 2007. Þrándheimur: SINTEF.
- Bollini, P., Tibaldi, G., Testa, C., og Munizza, C. (2004). Understanding treatment adherence in affective disorders: A qualitative study. *Journal of Psychiatric and Mental Health Nursing*, 11(6), 668-674.
- Bosc, M. (2000). Assessment of social functioning in depression. *Comprehensive Psychiatry*, 41(1), 63-69.
- Boschloo, L., Reeuwijk, K. G., Schoevers, R. A., og W J H Penninx, B. (2014). The impact of lifestyle factors on the 2-year course of depressive and/or anxiety disorders. *Journal of Affective Disorders*, 159, 73-79. doi:<http://dx.doi.org/10.1016/j.jad.2014.01.019>
- Bostwick, J. M., og Pankratz, V. S. (2000). Affective disorders and suicide risk: A reexamination. *The American Journal of Psychiatry*, 157(12), 1925-32
- Boyd, J. E., Otilingam, P. G., og DeForge, B. R. (2014). Brief version of the internalized stigma of mental illness (ISMI) scale: Psychometric properties and relationship to depression, self esteem, recovery orientation, empowerment, and perceived devaluation and discrimination. *Psychiatric Rehabilitation Journal*, 37(1), 17-23.
- Brodaty, H., Luscombe, G., Peisah, C., Anstey, K., og Andrews, G. (2001). A 25-year longitudinal, comparison study of the outcome of depression. *Psychological Medicine*, 31(8), 1347-59.

- Broocks, A., Meyer, T. F., Bandelow, B., George, A., Bartmann, U., Rüther, E., og Hillmer-Vogel, U. (1997). Exercise avoidance and impaired endurance capacity in patients with panic disorder. *Neuropsychobiology*, 36(4), 182-187.
- Brunello, N., Armitage, R., Feinberg, I., Holsboer-Trachsler, E., Léger, D., Linkowski, P., ... Mendlewicz, J. (2000). Depression and sleep disorders: Clinical relevance, economic burden and pharmacological treatment. *Neuropsychobiology*, 42(3), 107-19.
- Busch, K. A., Fawcett, J., og Jacobs, D. G. (2003). Clinical correlates of inpatient suicide. *Journal of Clinical Psychiatry*, 64(1), 14-19.
- Buttigieg, S. C., West, M. A., og Dawson, J. F. (2011). Well-structured teams and the buffering of hospital employees from stress. *Health Services Management Research*, 24(4), 203-212. doi:<http://dx.doi.org/10.1258/hsmr.2011.011013>
- Capdeville, D. og Ritchie, K. (2008). The long and the short of it: Are shorter periods of hospitalisation beneficial?, *The British Journal of Psychiatry*, 192(3), 164-165. doi: 10.1192/bjp.bp.107.044719
- Carrera, M., Herrán, A., Ayuso-Mateos, J. L., Sierra-Biddle, D., Ramírez, M. L., Ayestarán, A., ... Vázquez-Barquero, J. L. (2006). Quality of life in early phases of panic disorder: Predictive factors. *Journal of Affective Disorders*, 94(1-3), 127-134.
- Cassano, P. og Fava, M. (2002). Depression and public health an overview. *Journal of Psychosomatic Research*, 53, 849-857.
- Cavanagh, J. T. O., Carson, A. J., Sharpe, M., og Lawrie, S. M. (2003). Psychological autopsy studies of suicide: A systematic review. *Psychological Medicine*, 33(3), 395-405.
- Cepoiu, M., Mccusker, J., Cole, M. G., Sewitch, M., Belzile, E., og Ciampi, A. (2008). Recognition of depression by non-psychiatric physicians: A systematic literature review and meta-analysis. *Journal of General Internal Medicine*, 23(1), 25-36.
- Chan, M. F. (2007). Effects of music on patients undergoing a C-clamp procedure after percutaneous coronary interventions: A randomized controlled trial. *Heart & Lung : The Journal of Critical Care*, 36(6), 431-439.
- Chan, M. F., Chan, E. A., Mok, E., og Kwan Tse, F. Y. (2009). Effect of music on depression levels and physiological responses in community-based older adults. *International Journal of Mental Health Nursing*, 18(4), 285-294. doi:<http://dx.doi.org/10.1111/j.1447-0349.2009.00614.x>
- Chan, M.F., Wong, Z.Y. og Thayala, N.V. (2010). A systematic review of the effectiveness of music listening in reducing depressive symptoms in adults. *JBI Library of Systematic Reviews*, 8(31), 1242-1287.
- Chen, Y. og Li, I. (2009). Effectiveness of intervention using empowerment concept for patients with chronic disease: A systematic review. *JBI Library of Systematic Reviews*, 7(27), 1177-1232.

- Christer Magnusson. (2009a). Fjölskylduhjúkrun á Landspítala. *Tímarit hjúkrunarfræðinga*, 85(5), 12-16.
- Christer Magnusson. (2009b). Rannsakar fjölskylduhjúkrun á geðsviði. *Tímarit hjúkrunarfræðinga*, 85(5), 18-19.
- Cohen, S. og Herber, T.B. (1996). Health Psychology: Psychological factors and physical disease from the perspective of human psychoneuroimmunology, *Annual Review of Psychology*, 47, 113-142.
- Collins, S. (2005). Explanations in consultations: The combined effectiveness of doctors' and nurses' communication with patients. *Medical Education*, 39(8), 785-796.
- Connolly, N.P., Eberhart, N.K., Hammen, C.L. og Brennan, P.A. (2010). Specificity of Stress Generation: A Comparison of Adolescents. *Int J Cogn Ther*, 3(4), 368-379.
doi:10.1521/ijct.2010.3.4.368
- Cooper-patrick, L., Powe, N.R., Jenckes, M.W., Gonzales, J.J., Levine, D.M. og Ford, D.E. (1997). Identification of patient attitudes and preferences regarding treatment of depression. *Journal of General Internal Medicine*, 12(7), 431-8. doi:<http://dx.doi.org/10.1046/j.1525-1497.1997.00075.x>
- Coster, S. og Norman, I. (2009). Cochrane reviews of educational and self-management interventions to guide nursing practice: A review. *International Journal of Nursing Studies*, 46(4), 508-528.
doi: 10.1016/j.ijnurstu.2008.09.009
- Crowe, M., Beaglehole, B., Wells, H. og Porter, R. (2015). Non-pharmacological strategies for treatment of inpatient depression. *Australian and New Zealand Journal of Psychiatry*, 49(3), 215-226.
- Crowe, M., og Luty, S. (2005). Interpersonal psychotherapy: An effective psychotherapeutic intervention for mental health nursing practice. *International Journal of Mental Health Nursing*, 14(1), 24-31.
- Cuijpers, P., Berking, M., Anderson, G., Quigley, L., Kleiboer, A. og Dobson, K.S. (2013). A meta-analysis of cognitive-behavioural therapy for adult depression, alone and in comparison with other treatments, *Can J Psychiatry*, 58(7), 376-385.
- Currid, T.J. (2012). Cognitive behavioural group therapy is moderately effective for depression, with continued effect for up to 6 months but not beyond. *Evidence based nursing*, 9, 2-17.
- Demyttenaere, K. (2006). Quality of life in depression and anxiety: Does it matter? *International Journal of Psychiatry in Clinical Practice*, 10(1), 27-30.
- Department of mental health. (1999). *Mental health: A report of the surgeon General*. Rockville: National Institute of mental health.
- Derubeis, R.J., Hollon, S.D., Amsterdam, J.D., Shelton, R.C., Young, P.R., Salomon, R.M ... Gallop, R. (2005). Cognitive therapy vs medication in the treatment of moderate to severe depression. *Arch Gen Psychiatry*, 62(4), 409-416.

- Desplenter, F. A., Laekeman, G. J., og Simoens, S. R. (2011). Following up patients with depression after hospital discharge: A mixed methods approach. *International Journal of Mental Health Systems*, 5, 28-38.
- Diego, M. A., Jones, N. A., Field, T., Hernandez-Reif, M., Schanberg, S., Kuhn, C., . . . Galamaga, M. (1998). Aromatherapy positively affects mood, EEG patterns of alertness and math computations. *The International Journal of Neuroscience*, 96(3-4), 217-224.
- Dierckx, B., Heijnen, W. T., van den Broek, W. W., og Birkenhäger, T. K. (2012). Efficacy of electroconvulsive therapy in bipolar versus unipolar major depression: A meta-analysis. *Bipolar Disorders*, 14(2), 146-150. doi:<http://dx.doi.org/10.1111/j.1399-5618.2012.00997.x>
- Donahue, A. B. (2000). Riding the mental health pendulum: Mixed messages in the era of neurobiology and self-help movements. *Social Work*, 45(5), 427-38.
- Drapalski, A. L., Lucksted, A., Perrin, P. B., Aakre, J. M., Brown, C. H., DeForge, B. R., og Boyd, J. E. (2013). A model of internalized stigma and its effects on people with mental illness. *Psychiatric Services*, 64(3), 264-9.
- Drevets, W. C., Price, J. L., og Furey, M. L. (2008). Brain structural and functional abnormalities in mood disorders: Implications for neurocircuitry models of depression. *Brain Structure and Function*, 213(1-2), 93-118. doi:<http://dx.doi.org/10.1007/s00429-008-0189-x>
- Dunlop, B. W., Scheinberg, K., og Dunlop, A. L. (2013). Ten ways to improve the treatment of depression and anxiety in adults. *Mental Health in Family Medicine*, 10(3), 175-181.
- Eells, K. (2014). The use of music and singing to help manage anxiety in older adults. *Mental Health Practice*, 17(5), 10-17. doi:<http://dx.doi.org/10.7748/mhp2014.02.17.5.10.e861>
- Ebmeier, K. P., Donaghey, C., og Steele, J. D. (2006). Recent developments and current controversies in depression. *The Lancet*, 367(9505), 153-67.
- Eby, G. A., og Eby, K. L. (2006). Rapid recovery from major depression using magnesium treatment. *Medical Hypotheses*, 67(2), 362-370.
- Ellen, S., Selzer, R., Norman, T., og Blashki, G. (2007). Depression and anxiety: Pharmacological treatment in general practice. *Australian Family Physician*, 36(4), 222-228.
- Erikson, S. og Gard, G. (2011). Physical exercise and depression. *Physical Therapy Reviews*, 16(4), 261-268.
- ESEMeD/MHEDEA 2000 Investigators. (2004). Prevalence of mental disorders in Europe: Results from the European study of the epidemiology of mental disorders (ESEMeD) project. *Acta Psychiatrica Scandinavica*, 420, 21-27.
- Farajzadegan, Z., Koosha, P., Sufi, G.J. og Keshvari, M. (2013). The relationship between family function and women's well-being. *Iran J Nurs Midwifery Res*, 18(1), 9-13.
- Fava, M. og Kendler, K.S. (2000). Major Depressive Disorder. *Neuron*, 28, 335-341.

- Firth-Cozens, J. (2001). Cultures for improving patient safety through learning: The role of teamwork. *Quality in Health Care*, 10(2), 26-31.
- Fortinash, K.M. og Worret, P.A.H. (2003). *Psychiatric Nursing Care Plans*. St.Louis: Mosby.
- Fröjd, K., Håkansson, A., Karlsson, I., og Molarius, A. (2003). Deceased, disabled or depressed--a population-based 6-year follow-up study of elderly people with depression. *Social Psychiatry and Psychiatric Epidemiology*, 38(10), 557-62.
- Funnel, M.M. og Anderson, R.M. (2004). Empowerment and self-management of diabetes. *Clinical Diabetes*, 22(3), 123-127. doi: 10.2337/diaclin.22.3.123
- Furukawa, T. A., Yoshimura, R., Harai, H., Imaizumi, T., Takeuchi, H., Kitamura, T., og Takahashi, K. (2009). How many well vs. unwell days can you expect over 10 years, once you become depressed? *Acta Psychiatrica Scandinavica*, 119(4), 290-297.
- Garner, B., Phillips, L. J., Schmidt, H., Markulev, C., O'Connor, J., Wood, S. J., . . . McGorry, P. D. (2008). Pilot study evaluating the effect of massage therapy on stress, anxiety and aggression in a young adult psychiatric inpatient unit. *The Australian and New Zealand Journal of Psychiatry*, 42(5), 414-422. doi:<http://dx.doi.org/10.1080/00048670801961131>
- Gelder, M., Mayou, R. og Geddes, J. (1999). *Psychiatry*. New York: Oxford.
- George, M. S. (2002). New methods of minimally invasive brain modulation as therapies in psychiatry: TMS, MST, VNS and DBS. *Chinese Medical Journal*; 65(8), 349-360.
- Glick, I.D., Sharfstein, S.S. og Schwartz, H.I. (2011). Inpatient psychiatric care in the 21st century: The need for reform. *Psychiatric Services*, 62(2), 206-209. doi: 10.1176/appi.ps.62.2.206
- Gould, M. S., Greenberg, T., Velting, D. M., og Shaffer, D. (2003). Youth suicide risk and preventive interventions: A review of the past 10 years. *Journal of the American Academy of Child and Adolescent Psychiatry*, 42(4), 386-405.
- Griffiths, R.D., Fernandez, R.S., Mostacchi, M.S. og Evans, V. (2004). Comparison of educational interventions for mental health consumers receiving psychotropic medication. *Int J Evid Based Healthc*, 2(1), 1-44.
- Grigsby, A. B., Anderson, R. J., Freedland, K. E., Clouse, R. E., og Lustman, P. J. (2002). Prevalence of anxiety in adults with diabetes: A systematic review. *Journal of Psychosomatic Research*, 53(6), 1053-1060.
- Guðrún Pálmadóttir. (2013). Matstæki í rannsóknum: Öflun gagna um færni, þáttöku og umhverfi fólks. Í Sigríður Halldórsdóttir (ritstjóri), *Handbók í aðferðarfræði rannsókna* (1. Útgáfa). Akureyri: Háskólinn á Akureyri.
- Guyatt G.H., Naylor D., Juniper E., Heyland D.K., Jaeschke R. og Cook. D.J.(1997). Users' Guides to the Medical Literature. XII. How to use articles about Health-Related Quality of Life. *JAMA*, 2277: 1232-7.

Hawthorne, G., Goldney, R., og Taylor, A. W. (2008). Depression prevalence: Is it really increasing? *The Australian and New Zealand Journal of Psychiatry*, 42(7), 606-616.
doi:<http://dx.doi.org/10.1080/00048670802119788>

Heilbrigðis- og Tryggingarmálaráðuneytið. (2001). *Heilbrigðisáætlun til ársins 2010*. Reykjavík: Heilbrigðis- og Tryggingarmálaráðuneytið.

Heilbrigðis- og Tryggingarmálaráðuneytið. (2005). *Evrópuvígírlýsing um geðheilbrigðismál. Leitað lausna að brýnum verkefnum*. Sótt af http://www.velferdarraduneyti.is/media/frettir/Evropsk_yfirlysing_um_gedheilbrigdismal.pdf

Helmes, E. og Wiancko, D.C. (2006). Effects of music in reducing disruptive behavior in a general hospital. *Journal of the American Psychiatric Nurses Association*, 12(1), 37-44.

Hermes, B., Deakin, K., Lee, K. og Robinson, S. (2009). SUICIDE RISK assessment: 6 steps to a better instrument. *Journal of Psychosocial Nursing & Mental Health Services*, 47(6), 44-9.

Hewitt, J., og Coffey, M. (2005). Therapeutic working relationships with people with schizophrenia: Literature review. *Journal of Advanced Nursing*, 52(5), 561-570.

Hirschfeld, R.M.A. (2001). The comorbidity of major depression and anxiety disorders: Recognition and management in primary care. *Primary Care Companion J Clin Psychiatry*, 3(6), 244-254.

Hölzel, L., Härter, M., Reese, C. og Kriston, L. (2011). Risk factors for chronic depression – A systematic review. *Journal of Affective Disorders*, 129, 1-13.

International Council of Nurses. (2002) *Nurses Always There for You: Caring for Families. Information and Action Tool Kit*. Genf: International Council of Nurses.

Irwin, S. A., og Iglewicz, A. (2010). Oral ketamine for the rapid treatment of depression and anxiety in patients receiving hospice care. *Journal of Palliative Medicine*, 13(7), 903-908.

Isacsson, G., Reutfors, J., Papadopoulos, F. C., Ösby, U., og Ahlner, J. (2010). Antidepressant medication prevents suicide in depression. *Acta Psychiatrica Scandinavica*, 122(6), 454-460.

Jakob Smári, Daníel Þór Ólason, Þórður Örn Arnarson og Jón Friðrik Sigurðsson. (2008). Mælitæki fyrir þunglyndi fullorðinna sem til eru í íslenskri gerð: Próffræðilegar upplýsingar og notagildi. *Sálfræðiritið*, 13, 147-169.

Joint Commission (2009), “*Patient safety goals*”, sótt af www.jointcommission.org/standards_information/npsgs.aspx

Judd, L. L., Akiskal, H. S., Maser, J. D., Zeller, P. J., Endicott, J., Coryell, W. ... Keller, M.B. (2005). A prospective 12-year study of subsyndromal and syndromal depressive symptoms in unipolar major depressive disorders. *Archives of General Psychiatry*, 55(8), 694-700.

Kam-shing Yip. (2005). A strengths perspective in working with an adolescent with depression. *Psychiatric Rehabilitation Journal*, 28(4), 362-9.

- Kayser, S., Bewernick, B. H., Grubert, C., Hadrysiewicz, B. L., Axmacher, N., og Schlaepfer, T. E. (2011). Antidepressant effects of magnetic seizure therapy and electroconvulsive therapy, in treatment-resistant depression. *Journal of Psychiatric Research*, 45(5), 569-576.
- Kendler, K. S., Gardner, C. O., Neale, M. C., og Prescott, C. A. (2001). Genetic risk factors for major depression in men and women: Similar or different heritabilities and same or partly distinct genes? *Psychological Medicine*, 31(4), 605-16.
- Kennedy, N., Abbott, R., og Paykel, E. S. (2003). Remission and recurrence of depression in the maintenance era: Long-term outcome in a cambridge cohort. *Psychological Medicine*, 33(5), 827-38.
- Kennedy S.H., Lam R.W., Nutt D.J., Thase M.E. (2007). *Prevalance, burden and diagnosis. Treating depression effectively: applying clinical guidelines*. London: Martin Dunitz.
- Kessler, R. C., Berglund, P., Demler, O., Jin, R., Koretz, D., Merikangas, K. R., . . . Wang, P. S. (2003). The epidemiology of major depressive disorder: Results from the national comorbidity survey replication (NCS-R). *The Journal of the American Medical Association*, 289(23), 3095-3105.
- Kessler, R. C., Berglund, P., Demler, O., Jin, R., Merikangas, K.R. og Walters, E. E. (2005). Lifetime prevalence and age-of-onset distributions of DSM-IV disorders in the national comorbidity survey replication. *Archives of General Psychiatry*, 62(6), 593-602.
- Kessler, R. C., Borges, G., og Walters, E. E. (1999). Prevalence of and risk factors for lifetime suicide attempts in the national comorbidity survey. *Archives of General Psychiatry*, 56(7), 617-626.
- Kessler, R. C., Zhao, S., Blazer, D. G., og Swartz, M. (1997). Prevalence, correlates, and course of minor depression and major depression in the national comorbidity survey. *Journal of Affective Disorders*, 45(1-2), 19-30.
- Knapen, J., Van de Vliet, P., Van Coppenolle, H., David, A., Peuskens, J., Knapen, K., og Pieters, G. (2003). The effectiveness of two psychomotor therapy programmes on physical fitness and physical self-concept in nonpsychotic psychiatric patients: A randomized controlled trial. *Clinical Rehabilitation*, 17(6), 637-647.
- Knapen, J., Vermeersch, J., Van Coppenolle, H., Cuykx, V. Pieters, G. og Peuskens, J. (2007). The physical self-concept in patients with depressive and anxiety disorders. *International Journal of Therapy and Rehabilitation*, 14(1), 31-36.
- Koivumaa-honkanen, H., Rissanen, T., Hintikka, J., Honkalampi, K., Haatainen, K., Tarja, S., og Viinamäki, H. (2011). Factors associated with life satisfaction in a 6-year follow-up of depressive out-patients. *Soc Psychiatry Psychiatr Epidemiol*, 46, 595-605.
- Korkeila, J., Oksanen, T., Virtanen, M., Salo, P., Nabi, H., Pentti, J., . . . Kivimäki, M. (2011). Early retirement from work among employees with a diagnosis of personality disorder compared to anxiety and depressive disorders. *European Psychiatry : The Journal of the Association of European Psychiatrists*, 26(1), 18-22. doi:<http://dx.doi.org/10.1016/j.eurpsy.2009.12.022>

Kristín S. Jensdóttir og Sigurður B. Stefánsson. (2010). *Raflækningar – Leiðbeiningar fyrir notendur [Bæklingur]*. Reykjavík: Landspítalinn.

Kruijshaar ME, Barendregt, J., Vos, T., de Graaf R, Spijker, J., og Andrews, G. (2005). Lifetime prevalence estimates of major depression: An indirect estimation method and a quantification of recall bias. *European Journal of Epidemiology*, 20(1), 103-11.

Kumlien, S., og Axelsson, K. (2000). The nursing care of stroke patients in nursing homes. Nurses' descriptions and experiences relating to cognition and mood. *Journal of Clinical Nursing*, 9(4), 489-497.

Köhler, S., Unger, T., Hoffman, S. og Steinacher, B. (2013). Acute and long-term treatment outcome in depressed inpatients with vs. Without anxious features: Results of a one-year follow-up study, *Jounarl of Affective Disorders*, 150, 1055-1061.

Köhler, S., Wiethoff, K., Ricken, R., Stamm, T., Baghai, T.C., Fisher, R. ...Adli, M. (2015). Characteristics and differences in treatment outcome of inpatients with chronic vs. Episodic major depressive disorders, *Journal of Affective Disorders*, 173, 126-133.

Landlæknisembættið. (2012). *Sjúkrahúsið á Akureyri (FSA)*. Úttekt á gæðum og öryggi þjónustu. Reykjavík: Landlæknisembættið.

Large, M., Sharma, S., Cannon, E., Ryan, C., og Nielssen, O. (2011). Risk factors for suicide within a year of discharge from psychiatric hospital: A systematic meta-analysis. *The Australian and New Zealand Journal of Psychiatry*, 45(8), 619-628.
doi:<http://dx.doi.org/10.3109/00048674.2011.590465>

Lerner, D., Adler, D. A., Chang, H., Lapitsky, L., Hood, M. Y., Perissinotto, C., . . . Rogers, W. H. (2004). Unemployment, job retention, and productivity loss among employees with depression. *Psychiatric Services*, 55(12), 1371-1378.

Levin, F. R., Mariani, J., Brooks, D. J., Pavlicova, M., Nunes, E. V., Agosti, V., . . . Carpenter, K. M. (2013). A randomized double-blind, placebo-controlled trial of venlafaxine-extended release for co-occurring cannabis dependence and depressive disorders. *Addiction*, 108(6), 1084-1094.
doi:<http://dx.doi.org/10.1111/add.12108>

Lewinsohn, P. M., Rohde, P., Seeley, J. R., Klein, D. N., og Gotlib, I. H. (2003). Psychosocial functioning of young adults who have experienced and recovered from major depressive disorder during adolescence. *Journal of Abnormal Psychology*, 112(3), 353-363.

Limosin, F., Loze, J., Zylberman-Bouhassira, M., Schmidt, M.E., Perrin, E. og Rouillon, F. (2004). The course of depressive illness in general practice. *Can J Psychiatry*, 49, 119-123.

Lingam, R. og Scott, J. (2002). Treatment non-adherence in affective disorders. *Acta Psychiatr Scand*, 105, 164-172.

Lisanby, S. H., MD. (2007). Electroconvulsive therapy for depression. *The New England Journal of Medicine*, 357(19), 1939-45. doi:<http://dx.doi.org/10.1056/NEJMct075234>

- Liu, X., Gentzler, A. L., Tepper, P., Kiss, E., Kothencné, V. O., Tamás, Z., . . Kovacs, M. (2006). Clinical features of depressed children and adolescents with various forms of suicidality. *The Journal of Clinical Psychiatry*, 67(9), 1442-1450.
- Livingston, J. D., og Boyd, J. E. (2010). Correlates and consequences of internalized stigma for people living with mental illness: A systematic review and meta-analysis. *Social Science & Medicine*, 71(12), 2150-2161.
- Lloyd-Williams, M., Friedman, T., og Rudd, N. (1999). A survey of antidepressant prescribing in the terminally ill. *Palliative Medicine*, 13(3), 243-248.
doi:<http://dx.doi.org/10.1191/026921699676753309>
- Lopez, A. D., Mathers, C. D., Ezzati, M., Jamison, D. T., og Murray, C. J. L. (2006). Global and regional burden of disease and risk factors, 2001: Systematic analysis of population health data. *The Lancet*, 367(9524), 1747-57.
- Lorei, T.W. (1970). Staff ratings of the relative importance of the consequences of release from or retention in a psychiatric hospital. *Journal of Consulting and clinical psychology*, 34(1), 48-55.
- Lorenzetti, V., Allen, N. B., Fornito, A., og Yücel, M. (2009). Structural brain abnormalities in major depressive disorder: A selective review of recent MRI studies. *Journal of Affective Disorders*, 117(1-2), 1-17. doi:<http://dx.doi.org/10.1016/j.jad.2008.11.021>
- Lovibond, P.F. og Lovibond, S.H. (1995). The structure of negative emotional states: Comparison of the depression anxiety stress scales (DASS) with the beck depression and anxiety inventories. *Behav. Res. Ther.*, 33(3), 335-343.
- Lucksted, A., Drapalski, A., Calmes, C., Forbes, C., DeForge, B., og Boyd, J. (2011). Ending self-stigma: Pilot evaluation of a new intervention to reduce internalized stigma among people with mental illnesses. *Psychiatric Rehabilitation Journal*, 35(1), 51-4.
- Lustman, P. J., Anderson, R. J., Freedland, K. E., de Groot, M. og Carney, R.M. (2000). Depression and poor glycemic control: A meta-analytic review of the literature. *Diabetes Care*, 23(7), 934-42.
- Lög um persónuvernd nr. 77/2000.
- Lög um réttindi sjúklinga nr. 74/1997.
- Löwe, B., Spitzer, R. L., Williams, J. B. W., Mussell, M., Schellberg, D., og Kroenke, K. (2008). Depression, anxiety and somatization in primary care: Syndrome overlap and functional impairment. *General Hospital Psychiatry*, 30(3), 191-199.
- MacPherson, H., og Schroer, S. (2007). Acupuncture as a complex intervention for depression: A consensus method to develop a standardised treatment protocol for a randomised controlled trial. *Complementary Therapies in Medicine*, 15(2), 92-100.
doi:<http://dx.doi.org/10.1016/j.ctim.2006.09.006>

- Maj, M., Veltro, F., Pirozzi, R., Lobrace, S., og Magliano, L. (1992). Pattern of recurrence of illness after recovery from an episode of major depression: A prospective study. *The American Journal of Psychiatry*, 149(6), 795-800.
- Manser, T. (2009). Teamwork and patient safety in dynamic domains of healthcare: A review of the literature. *Acta Anaesthesiologica Scandinavica*, 53(2), 143-151.
doi:<http://dx.doi.org/10.1111/j.1399-6576.2008.01717.x>
- Marsh, D.T. (1999). Serious mental illness: Opportunities for family practitioners. *The Family Journal: Counseling And Therapy For Couples And Families*, 7(4), 358-366.
- Meades, R. og Ayers, S. (2011). Anxiety measures validated in perinatal populations: A systematic review. *Journal of Affective Disorders*, 133, 1-15.
- Mellesdal, L., Mehlum, L., Wentzel-Larsen, T., Kroken, R. og Jørgensen A.H. (2010). Suicide risk and acute psychiatric readmissions: A prospective cohort study. *Psychiatric Services*, 61(1), 25-31.
- Miller, G.E., Chen, E. og Parker, K.J.(2011). Psychological stress in childhood and susceptibility to the chronic diseases of aging: Moving toward a model of behavioral and biological mechanisms. *Psychological Bulletin*, 137(6), 959-997.
- Mitchell, A. J., Vaze, A., og Rao, S. (2009). Clinical diagnosis of depression in primary care: A meta-analysis. *The Lancet*, 374(9690), 609-19.
- Moret, C., og Briley, M. (2011). The importance of norepinephrine in depression. *Neuropsychiatric Disease and Treatment*, 7, 9-13. doi:<http://dx.doi.org/10.2147/NDT.S19619>
- Morris, D. W., Trivedi, M. H., og Rush, A. J. (2008). Folate and unipolar depression. *Journal of Alternative and Complementary Medicine*, 14(3), 277-285.
doi:<http://dx.doi.org/10.1089/acm.2007.0663>
- Moussavi, S., Chatterji, S., Verdes, E., Tandon, A. Patel, V. og han Ustun, B. (2007). Depression, chronic diseases, and decrements in health: Results from the world health surveys. *The Lancet*, 370(9590), 851-8.
- Mueller, T.I., Leon, A.C., Keller, M.B., Solomon, D.A., Endicott, J., Coryell, W ... Maser, J.D. (1999). Recurrence after recovery from major depressive disorder during 15 years of observational follow-up. *Am J Psychiatry*, 156(7), 1000-1006.
- Murray, C. J. L., og Lopez, A. D. (1997). Alternative projections of mortality and disability by cause 1990-2020: Global burden of disease study. *Lancet*, 349(9064), 1498-1504.
- National Institute for Health og Clinical Excellence (NICE) (2010). *Depression. The treatment and management of depression in adults* (updated edition). London: The British Psychological Society & The Royal College of Psychiatrists.

- Nicholson, A., Kuper, H., og Hemingway, H. (2006). Depression as an aetiologic and prognostic factor in coronary heart disease: A meta-analysis of 6362 events among 146 538 participants in 54 observational studies. *European Heart Journal*, 27(23), 2763-74.
- Nilsson, U. (2008). The anxiety- and pain-reducing effects of music interventions: A systematic review. *AORN Journal*, 87(4), 780-807. doi:<http://dx.doi.org/10.1016/j.aorn.2007.09.013>
- Noble, L.M., Douglas, B.C. og Newman, S.P. (2001). What do patients expect of psychiatric services? A systematic and critical review of empirical studies. *Social Science og Medicine*, 52, 985-998.
- Nolen-Hoeksema, S. (2007). *Abnormal psychology*. New York: McCraw-Hill.
- Nolen-Hoeksema, S., Girgus, J. S., og Seligman, M. E. (1992). Predictors and consequences of childhood depressive symptoms: A 5-year longitudinal study. *Journal of Abnormal Psychology*, 101(3), 405-422.
- Oeland, A., Laessoe, U., Olesen, A. V., og Munk-Jørgensen, P. (2010). Impact of exercise on patients with depression and anxiety. *Nordic Journal of Psychiatry*, 64(3), 210-217. doi:<http://dx.doi.org/10.3109/08039480903511373>
- Olfson, M., Mojtabai, R., Sampson, N. A., Hwang, I., Druss, B., Wang, Philip S. ... Kessler, R. C. (2009). Dropout from outpatient mental health care in the united states. *Psychiatric Services*, 60(7), 898-907.
- O'Reardon, J. P., Solvason, H. B., Janicak, P. G., Sampson, S., Isenberg, K. E., Nahas, Z., ... Sackeim, H. A. (2007). Efficacy and safety of transcranial magnetic stimulation in the acute treatment of major depression: A multisite randomized controlled trial. *Biological Psychiatry*, 62(11), 1208-1216.
- Osborn, D. P. J. (2001). The poor physical health of people with mental illness. *Western Journal of Medicine*, 175(5), 329-32.
- Overton, S. L., og Medina, S. L. (2008). The stigma of mental illness. *Journal of Counseling and Development*, 86(2), 143-151.
- Pampallona, S., Bollini, P., Tibaldi, G., Kupelnick, B., og Munizza, C. (2002). Patient adherence in the treatment of depression. *The British Journal of Psychiatry*, 180, 104-109.
- Páll Biering, Guðbjörg Daníelsdóttir og Arndís Ósk Jónsdóttir. (2005). *Pjónustuparfir geðsjúkra og reynsla heirra af geðheilbrigðispjónustu á Íslandi*. Reykjavík: Rauði Kross Íslands og Geðhjálps.
- Peritogiannis, V., Stefanou, E., Damigos, D., og Mavreas, V. (2008). Admission rates of patients with borderline personality disorder in a psychiatric unit in a general hospital. *International Journal of Psychiatry in Clinical Practice*, 12(1), 78-80.
- Peplau, H. E. (1989). *Interpersonal Theory in Nursing Practice Selected works of Hildegard E. Peplau*. Í Anita Werner O'Toole and Sheila Rouslin Welt (ritstjórar). New York: Springer Publishing Company.

- Perreault, M., Tardif, H., Provencher, H., Paquin, G., Desmarais, J., og Pawliuk, N. (2005). The role of relatives in discharge planning from psychiatric hospitals: The perspective of patients and their relatives. *Psychiatric Quarterly*, 76(4), 297-315.
- Perry, N., og Perry, E. (2006). Aromatherapy in the management of psychiatric disorders: Clinical and neuropharmacological perspectives. *CNS Drugs*, 20(4), 257-280.
- Pickett-Schenk, S. A., Cook, J. A., og Laris, A. (2000). Journey of hope program outcomes. *Community Mental Health Journal*, 36(4), 413-24.
- Pinto-Meza, A., Fernandez, A., Bruffaerts, R., Alonso, J., Kovess, V., Graaf, R., . . . Haro, J. M. (2011). Dropping out of mental health treatment among patients with depression and anxiety by type of provider: Results of the European study of the epidemiology of mental disorders. *Social Psychiatry and Psychiatric Epidemiology*, 46(4), 273-280.
- Pirkola, S., Sohlman, B., og Wahlbeck, K. (2005). The characteristics of suicides within a week of discharge after psychiatric hospitalisation - a nationwide register study. *BMC Psychiatry*, 5, 32.
- Powell, V.B., Abreu, N., de Oliveira, I.R. og Sudak, D. (2008). Cognitive-behavioral therapy for depression. *Rev Bras Psiquiatr*, 30(2), 73-80.
- Preti, A., Rucci, P., Gigantesco, A., Santone, G., Picardi, A., Miglio, R ... Zecca, L. (2009). Patterns of care in patients discharged from acute psychiatric inpatient facilities: A national survey in Italy. *Soc Psychiatry Epidemiol*, 44(9). 767-776. doi: 10.1007/s00127-009-0498-2
- Radfar, M., Ahmadi, F., og Fallahi Khoshknab, M. (2014). Turbulent life: The experiences of the family members of patients suffering from depression. *Journal of Psychiatric and Mental Health Nursing*, 21(3), 249-256. doi:<http://dx.doi.org/10.1111/jpm.12077>
- Ragnheiður Harpa Arnardóttir (2013). Megindlegar rannsóknir: Gerð rannsóknaráætlunar og yfirlit yfir helsu rannsóknarsnið. Í Sigríður Halldórsdóttir (ritstjóri), *Handbók í aðferðarfraeði rannsókna* (1. Útgáfa). Akureyri: Háskólinn á Akureyri.
- Ramachandani, P. og Stein, A. (2003) The impact of parental psychiatric disorder on children. *British Medical Journal*, 327, 242–243.
- Rayner, L., Price, A., Evans, A., Valsraj, K., Hotopf, M., og Higginson, I. J. (2011). Antidepressants for the treatment of depression in palliative care: Systematic review and meta-analysis. *Palliative Medicine*, 25(1), 36-51. doi:<http://dx.doi.org/10.1177/0269216310380764>
- Rihmer, Z., Appleby, L., Rihmer, A. og Belso, N. (2000). Decreasing suicide in Hungary, *The British Journal of Psychiatry*, 177(84), 84-89.
- Ritsher, J. B., Otilingam, P. G., og Grajales, M. (2003). Internalized stigma of mental illness: Psychometric properties of a new measure. *Psychiatry Research*, 121(1), 31-49.
- Rose, M. S., Koshman, M. L., Spreng, S., og Sheldon, R. (1999). Statistical issues encountered in the comparison of health-related quality of life in diseased patients to published general population norms: Problems and solutions. *Journal of Clinical Epidemiology*, 52(5), 405-412.

Rush, A. J., Trivedi, M. H., Wisniewski, S. R., Nierenberg, A. A., Stewart, J.W. ... Rosenbaum, J.F. (2006). Acute and longer-term outcomes in depressed outpatients requiring one or several treatment steps: A STAR*D report. *The American Journal of Psychiatry*, 163(11), 1905-17.

Sani, G., Tondo, L., Koukopoulos, A., Reginaldi, D., Kotzalidis, G. D., Koukopoulos, A. E., . . . Tatarelli, R. (2011). Suicide in a large population of former psychiatric inpatients. *Psychiatry and Clinical Neurosciences*, 65(3), 286-295. doi:<http://dx.doi.org/10.1111/j.1440-1819.2011.02205.x>

Sarris, J., og Kavanagh, D. J. (2009). Kava and st. john's wort: Current evidence for use in mood and anxiety disorders. *Journal of Alternative and Complementary Medicine*, 15(8), 827-836. doi:<http://dx.doi.org/10.1089/acm.2009.0066>

Sartorius, N. (2002) Eines der letzten Hindernisse einer verbesserten psychiatrischen Versorgung: das Stigma psychischer Erkrankung [One of the last obstacles to better mental health care: the stigma of mental illness]. *Neuropsychiatrie*, 16, 5–10.

Schlaepfer, T. E., Frick, C., Zobel, A., Maier, W., Heuser, I., Bajbouj, M., . . . Hasdemir, M. (2008). Vagus nerve stimulation for depression: Efficacy and safety in a european study. *Psychological Medicine*, 38(5), 651-61.

Schuch, F. B., Vasconcelos-Moreno, M. P., Borowsky, C., Zimmermann, A., Rocha, N. S., og Fleck, M. P. (2015). Exercise and severe major depression: Effect on symptom severity and quality of life at discharge in an inpatient cohort. *Journal of Psychiatric Research*, 61, 25-32. doi:<http://dx.doi.org/10.1016/j.jpsychires.2014.11.005>

Schober, M. og Affara, F. (2001). *The Family Nurse: Frameworks for practice*. Genf: International Council of Nurses.

Schulze-Rauschenbach, S. C., Harms, U., Schlaepfer, T. E., Maier, W., Falkai, P., og Wagner, M. (2005). Distinctive neurocognitive effects of repetitive transcranial magnetic stimulation and electroconvulsive therapy in major depression. *The British Journal of Psychiatry*, 186, 410-416.

Scope, A., Booth, A., og Sutcliffe, P. (2012). Women's perceptions and experiences of group cognitive behaviour therapy and other group interventions for postnatal depression: A qualitative synthesis. *Journal of Advanced Nursing*, 68(9), 1909-1919. doi:<http://dx.doi.org/10.1111/j.1365-2648.2012.05954.x>

Shapiro, D., Cook, I. A., Davydov, D. M., Ottaviani, C., Leuchter, A. F., og Abrams, M. (2007). Yoga as a complementary treatment of depression: Effects of traits and moods on treatment outcome. *Evidence-Based Complementary and Alternative Medicine*, 4(4), 493-502. doi:<http://dx.doi.org/10.1093/ecam/nel114>

Shuk Kwan Tang, og Tse, M. Y. M. (2014). Aromatherapy: Does it help to relieve pain, depression, anxiety, and stress in community-dwelling older persons? *BioMed Research International*, doi:<http://dx.doi.org/10.1155/2014/430195>

Sigríður Halldórsdóttir. (2013). Inngangur að aðferðarfræði. Í Sigríður Halldórsdóttir (ritstjóri), *Handbók í aðferðarfræði rannsókna* (1. Útgáfa). Akureyri: Háskólinn á Akureyri.

Sigurður Kristinsson. (2013). Siðfræði rannsókna og siðanefndir. Í Sigríður Halldórsdóttir (ritstjóri), *Handbók í aðferðarfræði rannsókna*. (1. Útgáfa). Akureyri: Háskólinn á Akureyri.

Sigurður Thorlacius og Sigurjón B. Stefánsson. (2009). *Algengi örorku á Íslandi 1.desember 2009 og samanburður við árin 2002 og 2005*. Sótt af <http://www.tr.is/media/skjol-a-pdf/Algengi-ororku-2009-Skyrsla.pdf>

Sigurður Thorlacius, Sigurjón B. Stefánsson og Stefán Ólafsson. (2007). Algengi örorku á Íslandi 1. desember 2005. *Læknablaðið*, 93(1), 11-14.

Sjúkrahúsið á Akureyri. (2015). *Gæðahandbók Sjúkrahússins á Akureyri*. Akureyri: Sjúkrahúsið á Akureyri.

Skorpen, A., Anderssen, N., Oye, C., og Bjelland, A. K. (2009). Treatment regimes in a psychiatric hospital and apparent contradictions: Report from field work in a lock-up ward illustrated with a constructed case. *International Journal of Mental Health Nursing*, 18(6), 409-416.
doi:<http://dx.doi.org/10.1111/j.1447-0349.2009.00630.x>

Smith, M., Ingram, T. og Brighton, V. (2009). Detection and assessment of late-life anxiety. *Journal of Gerontological*, 35(7), 9-15.

Sólrun Óladóttir. (2012). *Skjólstæðingsmiðuð þjónusta. Próun matstækis og starfsemi á geðdeild. FSA*. Heilbrigðisvínsindasvið. Háskólinn á Akureyri. Stell: Akureyri. Sótt af <http://skemman.is/item/view/1946/12477>

Spitzer, R. L., Kroenke, K., og Williams, J. B. W. (1999). Validation and utility of a self-report version of PRIME-MD: The PHQ primary care study. *Jama*, 282(18), 1737-44.

Steele, A., Maruyama, N., og Galynker, I. (2010). Psychiatric symptoms in caregivers of patients with bipolar disorder: A review. *Journal of Affective Disorders*, 121(1-2), 10-21.
doi:<http://dx.doi.org/10.1016/j.jad.2009.04.020>

Steinkuller, A. og Rheineck, J.E. (2009). A review of evidence-based therapeutic intervention for bipolar disorder. *Journal of Mental Health Counseling*, 31(4), 338-350.

Stjernswärd, S., og Ostman, M. (2007). Depression, e-health and family support. what the internet offers the relatives of depressed persons. *Nordic Journal of Psychiatry*, 61(1), 12-18.

Stone, M., Laughren, T., Jones, M. L., Levenson, M., Holland, P. C., Hughes, A., . . . Rochester, G. (2009). Risk of suicidality in clinical trials of antidepressants in adults: Analysis of proprietary data submitted to US food and drug administration. *British Medical Journal*, 339(7718), 431.

Sullivan, P. F., De Geus, E. J. C., Willemse, G., James, M. R., Smit, J. H., Zandbelt, T., . . . Penninx, B. W. J. H. (2009). Genome-wide association for major depressive disorder: A possible role for the presynaptic protein piccolo. *Molecular Psychiatry*, 14(4), 359-75.
doi:<http://dx.doi.org/10.1038/mp.2008.125>

- Sullivan, P. F., Neale, M. C., og Kendler, K. S. (2000). Genetic epidemiology of major depression: Review and meta-analysis. *The American Journal of Psychiatry*, 157(10), 1552-62.
- Svavarsdottir, E. K., Sigurdardottir, A. O., Konradsdottir, E., Stefansdottir, A., Sveinbjarnardottir, E. K., Ketilsdottir, A. . . . Guðmundsdottir, H. (2015). The process of translating family nursing knowledge into clinical practice. *Journal of Nursing Scholarship*, 47(1), 5-15.
- Sveinbjarnardottir, E. K., Svavarsdottir, E. K., og Hrafnkelsson, B. (2012). Psychometric development of the iceland-family perceived support questionnaire (ICE-FPSQ). *Journal of Family Nursing*, 18(3), 328-352. doi:<http://dx.doi.org/10.1177/1074840712449203>
- Sveinbjarnardottir, E. K., Svavarsdottir, E. K., og Saveman, B. (2011). Nurses attitudes towards the importance of families in psychiatric care following an educational and training intervention program. *Journal of Psychiatric and Mental Health Nursing*, 18(10), 895-903. doi:<http://dx.doi.org/10.1111/j.1365-2850.2011.01744.x>
- Sveinbjarnardottir, E.K., Svavarsdottir, E.K. og Wright, L.M. (2013). What are the benefits of a short therapeutic conversation intervention with acute psychiatric patients and their families? A controlled before and after study. *International Journal of Nursing Studies*, 50, 593-602. doi: 10.1016/j.ijnurstu.2012.10.009
- Svindseth, M. F., Nøttestad, J. A., og Dahl, A. A. (2010). A study of outcome in patients treated at a psychiatric emergency unit. *Nordic Journal of Psychiatry*, 64(6), 363-371.
- Terman, M., og Terman, J. S. (2005). Light therapy for seasonal and nonseasonal depression: Efficacy, protocol, safety, and side effects. *CNS Spectrums*, 10(8), 647-63.
- Thibeault, C.A., Trudeau, K., d'Entremont, M. og Brown. T. (2010). Understanding the milieu experiences of patients on an acute inpatient psychiatric unit. *Archives of Psychiatric Nursing*, 24(4), 216-226. doi: 10.1016/j.apnu.2009.07.002
- Thomas, A. J., Kalaria, R. N., og O'Brien, J. T. (2004). Depression and vascular disease: What is the relationship? *Journal of Affective Disorders*, 79(1-3), 81-95.
- Tiihonen, J., Lönnqvist, J., Wahlbeck, K., Klaukka, T., Tanskanen, A. og Haukka, J. (2006). Antidepressants and the risk of suicide, attempted suicide, and overall mortality in a nationwide cohort. *Archives of General Psychiatry*, 63(12), 1358-67.
- Trivedi, M.H. (2004). The link between depression and physical symptoms. *Prim Care Companion J Clin Psychiatry*, 6(1), 12–16.
- Trompenaars, F. J., Masthoff, E. D., Van Heck, G. L., Hodiamont, P. P., og De Vries, J. (2006). Relationship between mood related disorders and quality of life in a population of dutch adult psychiatric outpatients. *Depression and Anxiety*, 23(6), 353-363.
- Tómas Helgason. (1994). Faraldsfræðilegar rannsóknir í geðlæknið fræði á Íslandi. *Læknablaðið*, 80: 155-164.

Tómas Helgason, Helgi Tómasson og Tómas Zoëga (2004). Antidepressants and public health in Iceland. *British Journal of Psychiatry*, 184, 157-162.

Tómas Helgason, Júlíus K. Björnsson, Kristinn Tómasson og Erla Grétarsdóttir. (2000a). Heilsutengd lífsgæði Íslendinga. *Læknablaðið*, 86, 251-257.

Tómas Helgason, Júlíus K. Björnsson, Kristinn Tómasson, Erla Grétarsdóttir, Halldór Jónsson, Tómas Zoëga ... Guðmundur Vikar Einarsson (2000b) Heilsutengd lífsgæði sjúklinga fyrir og eftir meðferð. *Læknablaðið*, 86, 682-688.

Tómas Zoëga. (1984). Raflækningar. *Læknablaðið*, 70(10), 333-334.

Ustün, T. B., Ayuso-Mateos, J. L., Chatterji, S., Mathers, C., og Murray, C. J. L. (2004). Global burden of depressive disorders in the year 2000. *The British Journal of Psychiatry*, 184, 386-392.

van Beljouw, I. M., Verhaak, P. F., Cuijpers, P., van Marwijk, H. W., og Penninx, B. W. (2010). The course of untreated anxiety and depression, and determinants of poor one-year outcome: A one-year cohort study. *BMC Psychiatry*, 10, 86.

van Servellen, G., Heise, B. A., og Ellis, R. (2011). Factors associated with antidepressant medication adherence and adherence-enhancement programmes: A systematic literature review. *Mental Health in Family Medicine*, 8(4), 255-271.

Voelz, Z. R., Haefel, G. J., Joiner, T. E., og Dineen Wagner, K. (2003). Reducing hopelessness: The interaction of enhancing and depressogenic attributional styles for positive and negative life events among youth psychiatric inpatients. *Behaviour Research and Therapy*, 41(10), 1183-1198.

Vos, T., Corry, J., Haby, M. M., Carter, R., og Andrews, G. (2005). Cost-effectiveness of cognitive-behavioural therapy and drug interventions for major depression. *The Australian and New Zealand Journal of Psychiatry*, 39(8), 683-692.

Vuorilehto, M., Melartin, T., og Isometsä, E. (2005). Depressive disorders in primary care: Recurrent, chronic, and co-morbid. *Psychological Medicine*, 35(5), 673-82.

Wallsten, T., Kjellin, L. og Lindström, L. (2006). Short-term outcome of inpatient psychiatric care – impact of coercion and treatment characteristics. *Soc Psychiatry Epidemiol*, 41, 975-980.

Warren, B. J., og Lutz, W. J. (2007). The state of nursing science - cultural and lifespan issues in depression: Part I: Focus on adults. Issues in *Mental Health Nursing*, 28(7), 707-748.

Weber, M. (2010). The importance of exercise for individuals with chronic mental illness. *Journal of Psychosocial Nursing & Mental Health Services*, 48(10), 35-40.
doi:<http://dx.doi.org/10.3928/02793695-20100831-01>

Wells, K.B., Stewart, A., Hays, R.D., Burnam, A., Rogers, W., Daniels, M. ... Ware, J. (1989). The functioning and well-being of depressed patients: Results from the medical outcomes study. *JAMA*, 262, 914-919.

Wen-Hung, K., Gallo, J. J., og Eaton, W. W. (2004). Hopelessness, depression, substance disorder, and suicidality. *Social Psychiatry and Psychiatric Epidemiology*, 39(6), 497-501.
doi:<http://dx.doi.org/10.1007/s00127-004-0775-z>

West, M.A., Dawson, J.F., Admasachew, L. and Topakas, A. (2011), “NHS staff management And health service quality: results from the NHS Staff Survey and related data”. Sótt af www.dh.gov.uk/health/2011/08/nhs-staff-management/

West, M. A., og Lyubovnikova, J. (2013). Illusions of team working in health care. *Journal of Health Organization and Management*, 27(1), 134-142.

Wichers, M., Simons, C. J. P., Kramer, I. M. A., Hartmann, J. A., Lothmann, C., Myin-Germeys, I., . . . van Os, J. (2011). Momentary assessment technology as a tool to help patients with depression help themselves. *Acta Psychiatrica Scandinavica*, 124(4), 262-272.

World Health Organisation. (1995). The world health organization quality of life assessment (WHOQOL): Position paper from the world health organization. *Soc. Sci. Med*, 41(10), 1403-1409.

World Health Organisation. (2005). *Mental health: Facing the challenges, building solutions.* Kaupmannahöfn: World Health Organisation.

World Health Organisation. (2008). *The global burden of disease: 2004 update.* Genf: World Health Organisation.

World Health Organisation. (2012). *International Statistical Classification of Diseases and Related Health Problems 10th Revision.* Sótt af <http://apps.who.int/classifications/icd10/browse/2010/en>

World Health Organisation. (2014a). *Global status report on alcohol and health 2014.* Luxemborg. World Health Organisation.

World Health Organisation. (2014b). *Preventing suicide. A global imperative.* Luxemborg: World Health Organisation.

World Health Organisation. (e.d.). *Mental Health.* Sótt af <http://www.euro.who.int/en/what-we-do/health-topics/noncommunicable-diseases/mental-health>

Wright, L.M. og Leahey, M. (2011) (Upphaflega gefin út 2009) *.Hjúkrun og fjölskyldur – Leiðbeiningar um mat á fjölskyldum og meðferð.* (Anna María Hilmarsdóttir og Þórunn M. Lárusdóttir þýddu). Reykjavík: Háskólaútgáfan.

Yang, W., Zilov, A., Soewondo, P., Bech, O.M., Sekkal, F. og Home, P.D. (2010). Observational studies: Going beyond the boundaries of randomized controlled trials. *Diabetes Research and Clinical Practice*, 88(1), 3-9.

Yerevanian, B. I., Koek, R. J., Feusner, J. D., Hwang, S., og Mintz, J. (2004). Antidepressants and suicidal behaviour in unipolar depression. *Acta Psychiatrica Scandinavica*, 110(6), 452-458.

Yoshimasu, K., Kiyohara, C., og Miyashita, K. (2008). Suicidal risk factors and completed suicide: Meta-analyses based on psychological autopsy studies. *Environmental Health and Preventive Medicine*, 13(5), 243-256. doi:<http://dx.doi.org/10.1007/s12199-008-0037-x>

Zanardi, R., Barbini, B., Rossini, D., Bernasconi, A., Fregni, F., Padberg, F., . . . Colombo, C. (2008). New perspectives on techniques for the clinical psychiatrist: Brain stimulation, chronobiology and psychiatric brain imaging. *Psychiatry and Clinical Neurosciences*, 62(6), 627-637.

Zimmerman, M., McGlinchey, J. B., Young, D., og Chelminski, I. (2006). Diagnosing major depressive disorder introduction: An examination of the DSM-IV diagnostic criteria. *Journal of Nervous and Mental Disease*, 194(3), 151-154.