

References

- Adams, T., Black, K., MacIntyre, C., MacIntyre, I., & Dean, R. (2012). Connectivity modelling and network analysis of sea lice infection in Loch Fyne, west coast of Scotland. *Aquaculture Environment Interactions*, 3(1), 51–63. doi:10.3354/aei00052
- Allison, A. (2012). Organic Accumulation under Salmon Aquaculture Cages in Fossfjördur, Iceland (Master's). University of Akureyri.
- Amundrud, T. L., & Murray, A. G. (2009). Modelling sea lice dispersion under varying environmental forcing in a Scottish sea loch. *Journal of Fish Diseases*, 32(1), 27–44. doi:10.1111/j.1365-2761.2008.00980.x
- Anderson, E. J., McGillis, W. R., & Grosenbaugh, M. A. (2001). The boundary layer of swimming fish. *Journal of Experimental Biology*, 204(1), 81–102.
- Anon (2009) Status for norske laksebestander i 2009 og råd om beskatning (The status of Norwegian salmon stocks in 2009 and catch advice). 226 pp. *Report from the Scientific Advisory Committee for Atlantic Salmon Management in Norway* 110.1, Trondheim.
- Arnarlax,. (2014). Current investigation in Haganes (pp. 4-29).
- Asplin, L., Boxaspen, K., & Sandvik, A. D. (2011). Modeling the distribution and abundance of planktonic larval stages of *Lepeophtheirus salmonis* in Norway. In S. Jones & R. Beamish (Eds.), *Salmon Lice* (pp. 29–50). Oxford, UK: Wiley-Blackwell.
- Asplin, L., Johnsen, I. A., Sandvik, A. D., Albretsen, J., Sundfjord, V., Aure, J., & Boxaspen, K. K. (2014). Dispersion of salmon lice in the Hardangerfjord. *Marine Biology Research*, 10(3), 216–225. doi:10.1080/17451000.2013.810755
- Atlantic Salmon Federation. (n.d.). Salmon life cycle. Retrieved from <http://www.asf.ca/life-cycle.html>. Last accessed: 12.03.2015
- Backhaus, J. O. (1985). A three-dimensional model for the simulation of shelf sea dynamics. *Deutsche Hydrographische Zeitschrift*, 38(4), 165–187. doi:10.1007/BF02328975
- Bailey, R. J. E., Birkett, M. A., Ingvarsdóttir, A., Mordue A. J., Mordue, W., O'Shea, B., . . . Wadhams, L. J. (2006). The role of semiochemicals in host location and non-host avoidance by salmon louse (*Lepeophtheirus salmonis*) copepodids. *Canadian Journal of Fisheries and Aquatic Sciences*, 63(2), 448–456. doi:10.1139/f05-231
- Berg, O. K., & Berg, M. (1989). The duration of sea and freshwater residence of the sea trout, *Salmo trutta*, from the Vardnes River in northern Norway. *Environmental Biology of Fishes*, 24(1), 23–32. doi:10.1007/BF00001607
- Berland, B., & Margolis, L. (1983). The early history of 'Lakselus' and some nomenclatural questions relating to copepod parasites of salmon. *Sarsia*, 68(4), 281–288.
- Bjørn, P. A., & Finstad, B. (1997). The physiological effects of salmon lice infection on sea trout post smolts. *Nordic Journal of Freshwater Research*, 73, 60–72.
- Bjørn, P. A., & Finstad, B. (1998). The development of salmon lice (*Lepeophtheirus salmonis*) on artificially infected post smolts of sea trout (*Salmo trutta*). *Canadian Journal of Zoology*, 76(5), 970–977. doi:10.1139/z98-003
- Bjørn, P. A., & Finstad, B. (2002). Salmon lice, *Lepeophtheirus salmonis* (Krøyer), infestation in sympatric populations of Arctic char, *Salvelinus alpinus* (L.), and sea trout, *Salmo trutta* (L.), in areas near and distant from salmon farms. *ICES Journal of Marine Science*, 59(1), 131–139. doi:10.1006/jmsc.2001.1143

- Bjørn, P. A., Finstad, B., & Kristoffersen, R. (2001). Salmon lice infection of wild sea trout and Arctic charr in marine and freshwaters: the effects of salmon farms. *Aquaculture Research*, 32(12), 947–962. doi:10.1046/j.1365-2109.2001.00627.x
- Bjørn, P. A., Finstad, B., Kristoffersen, R., McKinley, R. S., & Rikardsen, A. H. (2007). Differences in risks and consequences of salmon louse, *Lepeophtheirus salmonis* (Kroyer), infestation on sympatric populations of Atlantic salmon, brown trout, and Arctic charr within northern fjords. *ICES Journal of Marine Science*, 64(2), 386–393. doi:10.1093/icesjms/fsl029
- Bjørn, P. A., Sivertsgård, R., Finstad, B., Nilsen, R., Serra-Llinares, R. M., & Kristoffersen, R. (2011). Area protection may reduce salmon louse infection risk to wild salmonids. *Aquaculture Environment Interactions*, 1(3), 233–244. doi:10.3354/aei00023
- Boxaspen, K. (2006). A review of the biology and genetics of sea lice. *ICES Journal of Marine Science*, 63(7), 1304–1316. doi:10.1016/j.icesjms.2006.04.017
- Boxaspen, K., & Næss, T. (2000). Development of eggs and the planktonic stages of salmon lice (*Lepeophtheirus salmonis*) at low temperatures. *Contributions to Zoology*, 69(1/2), 51–56.
- Brabrand, Å., Koestler, A. G., & Borgstrøm, R. (2002). Lake spawning of brown trout related to groundwater influx. *Journal of Fish Biology*, 60(3), 751–763. doi:10.1111/j.1095-8649.2002.tb01699.x
- Bricknell, I. R., Dalesman, S. J., O Shea, B., Pert, C. C., & Mordue Luntz, A. J. (2006). Effect of environmental salinity on sea lice *Lepeophtheirus salmonis* settlement success. *Diseases of aquatic organisms*, 71(3), 201.
- Bron, J. E., Sommerville, C., Wootten, R., & Rae, G. H. (1993). Fallowing of marine Atlantic salmon, *Salmo salar* L., farms as a method for the control of sea lice, *Lepeophtheirus salmonis* (Kroyer, 1837). *Journal of Fish Diseases*, 16(5), 487–493. doi:10.1111/j.1365-2761.1993.tb00882.x
- Bush, A. O., Lafferty, K. D., Lotz, J. M., & Shostak, A. W. (1997). Parasitology meets ecology on its own terms: Margolis et al. revisited. *The Journal of parasitology*, 575–583.
- Butler, J. R. A. (2002). Wild salmonids and sea louse infestations on the west coast of Scotland: sources of infection and implications for the management of marine salmon farms. *Pest Management Science*, 58(6), 595–608. doi:10.1002/ps.490
- Butler, J. R. A., Watt, J., & Mills, D. (2003). Assessing and managing the impacts of marine salmon farms on wild Atlantic salmon in western Scotland: identifying priority rivers for conservation. *Salmon at the edge*, 93–118.
- Campbell, J. S. (1977). Spawning characteristics of brown trout and sea trout *Salmo trutta* L. in Kirk Burn, River Tweed, Scotland. *Journal of Fish Biology*, 11(3), 217–229. doi:10.1111/j.1095-8649.1977.tb04115.x
- Carlsen, K. T., Berg, O. K., Finstad, B., & Heggberget, T. G. (2004). Diel periodicity and environmental influence on the smolt migration of Arctic charr, *Salvelinus alpinus*, Atlantic salmon, *Salmo salar*, and brown trout, *Salmo trutta*, in northern Norway. *Environmental Biology of Fishes*, 70(4), 403–413. doi:10.1023/B:EBFI.0000035438.85321.fa
- Connors, B., Krkošek, M., & Dill, L. (2008). Sea lice escape predation on their host. *Biology Letters*, 4(5), 455–457. doi:10.1098/rsbl.2008.0276
- Costello, M. J. (1993). Review of methods to control sea lice (Caligidae: Crustacea) infestations on salmon (*Salmo salar*) farms. *Pathogens of wild and farmed fish: sea lice*, 219–252.
- Costello, M. J. (2006). Ecology of sea lice parasitic on farmed and wild fish. *Trends in Parasitology*, 22(10), 475–483. doi:10.1016/j.pt.2006.08.006
- Costello, M. J. (2009a). How sea lice from salmon farms may cause wild salmonid declines in Europe and North America and be a threat to fishes elsewhere. *Proceedings of the Royal Society B: Biological Sciences*, 276(1672), 3385–3394. doi:10.1098/rspb.2009.0771
- Costello, M. J. (2009b). The global economic cost of sea lice to the salmonid farming industry. *Journal of Fish Diseases*, 32(1), 115–118. doi:10.1111/j.1365-2761.2008.01011.x
- Costelloe, J., Costelloe, M., & Roche, N. (1995). Variation in sea lice infestation on Atlantic salmon smolts in Killary Harbour, West Coast of Ireland. *Aquaculture International*, 3(4), 379–393. doi:10.1007/BF00121626

- D. Todd, C., Walker, A. M., Hoyle, J. E., Northcott, S. J., Walker, A. F., & Ritchie, M. G. (2000). Infestations of wild adult Atlantic salmon (*Salmo salar* L.) by the ectoparasitic copepod sea louse *Lepeophtheirus salmonis* Krøyer: prevalence, intensity and the spatial distribution of males and [2pt] females on the host fish. *Hydrobiologia*, 429(1/3), 181–196. doi:10.1023/A:1004031318505
- Davidsen, J. G., Plantalech Manel-la, N., Økland, F., Diserud, O. H., Thorstad, E. B., Finstad, B., . . . Rikardsen, A. H. (2008). Changes in swimming depths of Atlantic salmon *Salmo salar* post-smolts relative to light intensity. *Journal of Fish Biology*, 73(4), 1065–1074. doi:10.1111/j.1095-8649.2008.02004.x
- Dawson, L., Pike, A. W., Houlihan, D. F., & McVicar, A. H. (1997). Comparison of the susceptibility of sea trout (*Salmo trutta* L.) and Atlantic salmon (*Salmo salar* L.) to sea lice (*Lepeophtheirus salmonis* (Krøyer, 1837)) infections. *ICES Journal of Marine Science*, 54(6), 1129–1139. doi:10.1016/S1054-3139(97)80018-5
- Dempson, J. B., & Green, J. M. (1985). Life history of anadromous Arctic charr, *Salvelinus alpinus* in the Fraser River, northern Labrador. *Canadian Journal of Zoology*, 63(2), 315–324. doi:10.1139/z85-048
- Dempson, J. B., & Kristofferson A. H. (1987). Spatial and temporal aspects of the ocean migration of anadromous Arctic charr. *American Fisheries Society Symposium* (Vol. 1, pp. 340–357).,
- Dempson, J. B., O'Connell, M. F., & Shears, M. (1996). Relative production of Atlantic salmon from fluvial and lacustrine habitats estimated from analyses of scale characteristics. *Journal of Fish Biology*, 48(3), 329–341. doi:10.1111/j.1095-8649.1996.tb01431.x
- Ducharme, L. J. A. (1969). Atlantic salmon returning for their fifth and sixth consecutive spawning trips. *Journal of the Fisheries Research Board of Canada*, 26(6), 1661–1664. doi:10.1139/f69-149
- Dymond, J. R. (1963). Family salmonidae. *in-chief. Fishes of the Western North Atlantic. Sears Foundation Mar. Res.*, (1 Part 3), 457.
- Einarsson, S. M., Mills, D. H., & Johannsson, V. (1990). Utilisation of fluvial and lacustrine habitat by anadromous Atlantic salmon, *Salmo salar* L., in an Icelandic watershed. *Fisheries Research*, 10(1), 53–71. doi:10.1016/0165-7836(90)90015-N
- Elliott, J. M., & Hurley, M. A. (1998). An individual-based model for predicting the emergence period of sea trout fry in a lake district stream. *Journal of Fish Biology*, 53(2), 414–433. doi:10.1111/j.1095-8649.1998.tb00990.x
- Erkinaro, J., & Gibson, R. J. (1997). Interhabitat migration of juvenile Atlantic salmon in a Newfoundland river system, Canada. *Journal of Fish Biology*, 51(2), 373–388. doi:10.1111/j.1095-8649.1997.tb01673.x
- Euzenat, G., Fournel, F., & Richard, A. (2000). Sea trout (*Salmo trutta* L.) in Normandy and Picardy. In J. L. Baglinière & G. Maisse (Eds.), *Biology and Ecology of the Brown and Sea Trout* (pp. 175–203). London: Springer London.
- FAO. (2010). The state of world fisheries and aquaculture, 2010. Rome: *Food and Agriculture Organization of the United Nations*.
- FAO. (2014). State of World Fisheries and Aquaculture 2014: *Food and Agriculture Organization of the United Nations*.
- Fiksen, Ø., Jørgensen, C., Kristiansen, T., Vikebø, F., & Huse, G. (2007). Linking behavioural ecology and oceanography: larval behaviour determines growth, mortality and dispersal. *Marine Ecology Progress Series*, 347, 195–205. doi:10.3354/meps06978
- Finstad, B., Bjørn, P. A., Grimnes, A., & Hvidsten, N. A. (2000). Laboratory and field investigations of salmon lice [*Lepeophtheirus salmonis* (Kroyer)] infestation on Atlantic salmon (*Salmo salar* L.) post-smolts. *Aquaculture Research*, 31(11), 795–803. doi:10.1046/j.1365-2109.2000.00511.x
- Finstad, B., Bjørn, P. A., Todd, C. D., Whoriskey, F., Gargan, P. G., Forde, G., & Revie, C. W. (2011). The effect of sea lice on Atlantic salmon and other salmonid species. *Atlantic salmon ecology*, 253–276.
- Finstad, B., Økland, F., Thorstad, E. B., Bjørn, P. A., & McKinley, R. S. (2005). Migration of hatchery-reared Atlantic salmon and wild anadromous brown trout post-smolts in a Norwegian fjord system. *Journal of Fish Biology*, 66(1), 86–96. doi:10.1111/j.0022-1112.2005.00581.x

- Fjørtoft, H. B., Borgstrøm, R., & Skaala, Ø. (2013). Differential changes in growth patterns of anadromous brown trout and Atlantic salmon from the River Etneelva over a 25-year period. *Marine Biology Research*, 10(3), 301–307. doi:10.1080/17451000.2013.810753
- Ford, J. S., & Myers†, R. A. (2008). A global assessment of salmon aquaculture impacts on wild salmonids. *PLoS Biology*, 6(2), 411-417. doi:10.1371/journal.pbio.0060033
- Frazer, L. N. (2009). Sea-cage aquaculture, sea lice, and declines of wild fish. *Conservation Biology*, 23(3), 599–607. doi:10.1111/j.1523-1739.2008.01128.x
- Galbraith, M. (2005). Identification of larval stages of *Caligus clemensi* and *Lepeophtheirus salmonis* from the Broughton archipelago. *Canadian Technical Report of Fisheries and Aquatic Sciences* 2548:414
- Garcia-Vazquez, E., Moran, P., Martinez, J. L., Perez, J., Gaudemar, B. de, & Beall, E. (2001). Alternative mating strategies in Atlantic salmon and brown trout. *Journal of Heredity*, 92(2), 146–149. doi:10.1093/jhered/92.2.146
- Gargan, P. G., Poole, W. R., & Forde, G. P. (2006). A review of the status of Irish sea trout stocks. In G. Harris & N. Milner (Eds.), *Sea Trout: Biology, Conservation and Management* (pp. 25–44). Oxford, UK: Blackwell Publishing Ltd.
- Gargan, P. G., Tully, O., & Poole, W. R. (2003). The relationship between sea lice infestation, sea lice production and sea trout survival in Ireland, 1992–2001. *Salmon at the edge*, 119–135.
- Genna, R. L. (2002). The chemical ecology, physiology and infection dynamics of the sea louse copepodid, *Lepeophtheirus salmonis* kroyer. *UNIVERSITY OF ABERDEEN (UNITED KINGDOM)*.
- Genna, R. L., Mordue, W., Pike, A. W., & Mordue A. J. (2005). Light intensity, salinity, and host velocity influence presettlement intensity and distribution on hosts by copepodids of sea lice, *Lepeophtheirus salmonis*. *Canadian Journal of Fisheries and Aquatic Sciences*, 62(12), 2675–2682. doi:10.1139/f05-163
- Gillibrand, P. A., & Amundrud, T. L. (2007). A numerical study of the tidal circulation and buoyancy effects in a Scottish fjord: Loch Torridon. *Journal of Geophysical Research*, 112(C5). doi:10.1029/2006JC003806
- Gillibrand, P. A., & Willis, K. J. (2007). Dispersal of sea louse larvae from salmon farms: modelling the influence of environmental conditions and larval behaviour. *Aquatic Biology*, 1(1), 63–75.
- Gjelland, K. Ø., Serra-Llinares, R. M., Hedger, R. D., Arechavala-Lopez, P., Nilsen, R., Finstad, B., . . . Bjørn, P. A. (2014). Effects of salmon lice infection on the behaviour of sea trout in the marine phase. *Aquaculture Environment Interactions*, 5(3), 221–233. doi:10.3354/aei00105
- Grant, A. N. (2002). Medicines for sea lice. *Pest Management Science*, 58(6), 521–527. doi:10.1002/ps.481
- Grimnes, A., & Jakobsen, P. J. (1996). The physiological effects of salmon lice infection on post-smolt of Atlantic salmon. *Journal of Fish Biology*, 48(6), 1179–1194. doi:10.1111/j.1095-8649.1996.tb01813.x
- Guðbergsson, G. (2014). Catch statistics for Atlantic salmon Arctic charr and brown trout in Icelandic rivers and lakes. VMST/14045. Reykjavík: Institute of Freshwater Fisheries.
- Guðbergsson, G. & Antonsson, P. (1996). Fiskar í ám og vötnum – Fræðirit fyrir almenning um íslenska ferskvatnsfiska. Reykjavík: Landvernd.
- Guðjónsson, P. (1978). The Atlantic salmon in Iceland.
- Guðjónsson, P. (1990). Íslensk vötn og vistfræðileg flokkun þeirra. In: G. Sigbjarnarson (editor) Vatnið og landið, vatnafræðiráðstefna, október 1987 (pp.219-223). Reykjavík: Orkustofnun.
- Gulseth, O. A., & Nilssen, K. J. (2000). The brief period of spring migration, short marine residence, and high return rate of a northern Svalbard population of arctic char. *Transactions of the American Fisheries Society*, 129(3), 782–796.
- Haidvogel, D. B., Arango, H., Budgell, W. P., Cornuelle, B. D., Curchitser, E., Di Lorenzo, E., . . . Wilkin, J. (2008). Ocean forecasting in terrain-following coordinates: Formulation and skill assessment of the Regional Ocean Modeling System. *Journal of Computational Physics*, 227(7), 3595–3624. doi:10.1016/j.jcp.2007.06.016
- Halvorsen, M., & Svenning, M. (2000). Growth of Atlantic salmon parr in fluvial and lacustrine habitats. *Journal of Fish Biology*, 57(1), 145–160. doi:10.1111/j.1095-8649.2000.tb00782.x

- Hamre, L. A., Eichner, C., Caipang, Christopher Marlowe A., Dalvin, S. T., Bron, J. E., Nilsen, F., . . . Krkošek, M. (2013). The salmon louse *Lepeophtheirus salmonis* (Copepoda: Caligidae) life cycle has only two chalimus stages. *PLoS ONE*, 8(9), e73539. doi:10.1371/journal.pone.0073539
- Hayward, C. J., Andrews, M., & Nowak, B. F. (2011). Introduction: *Lepeophtheirus salmonis* -A remarkable success story. In S. Jones & R. Beamish (Eds.), *Salmon Lice* (pp. 1–28). Oxford, UK: Wiley-Blackwell.
- Hermansen, O. & Troell, M. (2012). *Aquaculture in the Arctic - a review* (Nofima Report 36/2012).
- Heuch, P. A. (1995). Experimental evidence for aggregation of salmon louse copepodids (*Lepeophtheirus salmonis*) in step salinity gradients. *Journal of the Marine Biological Association of the United Kingdom*, 75(04), 927. doi:10.1017/S002531540003825X
- Heuch, P. A., Bjørn, P. A., Finstad, B., Asplin, L., Holst, J. C. (2009) Salmon lice infection of farmed and wild salmonids in Norway: an overview. *Integr Comp Biol* 49(Suppl 1): e74, doi:10.1093/icb/icp002
- Heuch, P. A., Bjørn, P. A., Finstad, B., Holst, J. C., Asplin, L., & Nilsen, F. (2005). A review of the Norwegian ‘National Action Plan against Salmon Lice on Salmonids’: The effect on wild salmonids. *Aquaculture*, 246(1-4), 79–92. doi:10.1016/j.aquaculture.2004.12.027
- Heuch, P. A., Doall, M. H., & Yen, J. (2006). Water flow around a fish mimic attracts a parasitic and deters a planktonic copepod. *Journal of Plankton Research*, 29(Supplement 1), i3. doi:10.1093/plankt/fbl060
- Heuch, P. A., & Mo, T. A. (2001). A model of salmon louse production in Norway: effects of increasing salmon production and public management measures. *Diseases of aquatic organisms*, 45(2), 145–152.
- Heuch, P. A., Nordhagen, J. R., & Schram, T. A. (2000). Egg production in the salmon louse [*Lepeophtheirus salmonis* (Kroyer)] in relation to origin and water temperature. *Aquaculture Research*, 31(11), 805–814. doi:10.1046/j.1365-2109.2000.00512.x
- Heuch, P. A., Parsons, A., & Boxaspen, K. (1995). Diel vertical migration: A possible host-finding mechanism in salmon louse (*Lepeophtheirus salmonis*) copepodids? *Canadian Journal of Fisheries and Aquatic Sciences*, 52(4), 681–689. doi:10.1139/f95-069
- Hevroy, E., Boxaspen, K., Oppedal, F., Taranger, G., & Holm, J. (2003). The effect of artificial light treatment and depth on the infestation of the sea louse *Lepeophtheirus salmonis* on Atlantic salmon (*Salmo salar* L.) culture. *Aquaculture*, 220(1-4), 1–14. doi:10.1016/S0044-8486(02)00189-8
- Høgåsen, H. R. (1998). Physiological changes associated with the diadromous migration of salmonids (Vol. 127): *NRC Research Press*.
- Hvidsten, N. A., Johnsen, B. O. & Levings, C. D. 1992 Atferd og ernæring hos utvandrende laksesmolt i Trondheimsfjorden Behaviour and feeding of postsmolts of Atlantic salmon in the Trondheimsfjord). *NINA Oppdragsmelding*, 164, 14 (In Norwegian).
- Hvidsten, N. A., Johnsen, B. O. & Levings, C. D. 1995 Vandring og ernæring hos laksesmolt i Trondheims- fjorden og på Frohavet (Behaviour and feeding of postsmolts of Atlantic salmon in the Trondheimsfjord and Frohavet). *NINA Oppdragsmelding*, 332, 17 (In Norwegian).
- Ísaksson, Á. (2001). Major provisions in Icelandic laws and regulations. MONDARIZ, SPAIN. Retrieved from <http://www.mast.is/uploads/document/veidimal/utgafa/oslorespresl2001-001.pdf>
- Jackson, D., Deady, S., Leahy, Y., & Hassett, D. (1997). Variations in parasitic caligid infestations on farmed salmonids and implications for their management. *ICES Journal of Marine Science*, 54(6), 1104–1112. doi:10.1016/S1054-3139(97)80015-X
- Jackson, D., O'Donohoe, P., Kane, F., Kelly, S., Dermott, T. M., Drumm, A., . . . Nolan, G. (2012). Result of an epidemiological study of sea lice infestation in South Connemara, West of Ireland. *Aquaculture*, 364-365, 118–123. doi:10.1016/j.aquaculture.2012.08.003
- Jansen, P. A., Kristoffersen, A. B., Viljugrein, H., Jimenez, D., Aldrin, M., & Stien, A. (2012). Sea lice as a density-dependent constraint to salmonid farming. *Proceedings of the Royal Society B: Biological Sciences*, 279(1737), 2330–2338. doi:10.1098/rspb.2012.0084
- Järvi, T., Holmgren, K., Rubin, J. F., Petersson, E., Lundberg, S., & Glimsäter, C. (1996). Newly-emerged *Salmo trutta* fry that migrate to the sea-an alternative choice of feeding habit? *Nordic Journal of Freshwater Research* 72, 52–62.

- Jaworski, A., & Holm, J. C. (1992). Distribution and structure of the population of sea lice, *Lepeophtheirus salmonis* Krøyer, on Atlantic salmon, *Salmo salar* L., under typical rearing conditions. *Aquaculture Research*, 23(5), 577–589. doi:10.1111/j.1365-2109.1992.tb00802.x
- Jensen, A. J., Finstad, B., Fiske, P., Hvidsten, N. A., Rikardsen, A. H., Saksgård, L., & Bradford, M. (2012). Timing of smolt migration in sympatric populations of Atlantic salmon (*Salmo salar*), brown trout (*Salmo trutta*), and Arctic char (*Salvelinus alpinus*). *Canadian Journal of Fisheries and Aquatic Sciences*, 69(4), 711–723. doi:10.1139/f2012-005
- Jensen, K. W., & Berg, M. (1977). Growth, mortality and migrations of the anadromous char, *Salvelinus alpinus* L., in the Vardnes River, Troms, northern Norway. *Report Institute of Freshwater Research, Drottningholm*,
- Johnsen, I. A., Fiksen, Ø., Sandvik, A. D., & Asplin, L. (2014). Vertical salmon lice behaviour as a response to environmental conditions and its influence on regional dispersion in a fjord system. *Aquaculture Environment Interactions*, 5(2), 127–141. doi:10.3354/aei00098
- Johnson, L. (1989). The anadromous Arctic charr, *Salvelinus alpinus*, of Nauyuk Lake, NWT, Canada. *Physiology and Ecology Japan Special*, 1, 201–227.
- Johnson, S. C., & Albright, L. J. (1991a). Development, growth, and survival of *Lepeophtheirus salmonis* (Copepoda: Caligidae) under laboratory conditions. *Journal of the Marine Biological Association of the United Kingdom*, 71(02), 425. doi:10.1017/S0025315400051687
- Johnson, S. C., & Albright, L. J. (1991b). The developmental stages of *Lepeophtheirus salmonis* (Krøyer, 1837) (Copepoda: Caligidae). *Canadian Journal of Zoology*, 69(4), 929–950. doi:10.1139/z91-138
- Johnson, S. C., Blaylock, R. B., Elphick, J., & Hyatt, K. D. (1996). Disease induced by the sea louse ((*Lepeophtheirus salmonis*)) (Copepoda: Caligidae) in wild sockeye salmon (*Oncorhynchus nerka*) stocks of Alberni Inlet, British Columbia. *Canadian Journal of Fisheries and Aquatic Sciences*, 53(12), 2888–2897. doi:10.1139/f96-226
- Johnson, S. C., Fast, M. D., Wiegertjes, G., & Flik, G. (2004). Interactions between sea lice and their hosts. *Host-parasite interactions*, 131–159.
- Jones, J. W. (1959). The salmon: *Collins London*.
- Jones, M. W., Sommerville, C., & Bron, J. E. (1990). The histopathology associated with the juvenile stages of *Lepeophtheirus salmonis* on the Atlantic salmon, *Salmo salar* L. *Journal of Fish Diseases*, 13(4), 303–310. doi:10.1111/j.1365-2761.1990.tb00786.x
- Jones, S., & Beamish, R. (2011). Salmon lice: an integrated approach to understanding parasite abundance and distribution: *John Wiley & Sons*.
- Jónsson, G. (1983). Íslenskir fiskar. *Reykjavík: Fjölvautgáfan*.
- Jonsson, B., & Jonsson, N. (2009). Migratory timing, marine survival and growth of anadromous brown trout *Salmo trutta* in the River Imsa, Norway. *Journal of Fish Biology*, 74(3), 621–638. doi:10.1111/j.1095-8649.2008.02152.x
- Jonsson, B., & L'Abée-Lund, J. H. (1993). Latitudinal clines in life-history variables of anadromous brown trout in Europe. *Journal of Fish Biology*, 43(sa), 1–16. doi:10.1111/j.1095-8649.1993.tb01175.x
- Jonsson, G. S. (2000). Licensing, monitoring and regulation of aquaculture in Iceland. *Journal of Applied Ichthyology*, 16(4-5), 172–176. doi:10.1046/j.1439-0426.2000.00264.x
- Jørgensen, L., Amundsen, P.-A., Gabler, H.-M., Halvorsen, M., Erkinaro, J., & Niemelä, E. (1999). Spatial distribution of Atlantic salmon parr (*Salmo salar* L.) and bullhead (*Cottus gobio* L.) in lotic and lentic habitats of a diversified watercourse in northern Fennoscandia. *Fisheries Research*, 41(2), 201–211. doi:10.1016/S0165-7836(99)00014-4
- Kabata, Z. (1979). Parasitic copepoda of British fishes. [Publications] - Ray Society: no. 152. London: Ray Society.
- Karbowski, N. (2015). Assessment of sea lice infection rates on wild populations of salmonids in Arnarfjörður, Iceland (Masters). University of Akureyri.
- Kjartansdóttir, H. (2008). Repeat spawning of the Atlantic salmon (*Salmo salar*) in various salmon rivers in Iceland.

- Klemetsen, A., Amundsen, P.-A., Dempson, J. B., Jonsson, B., Jonsson, N., O'Connell, M. F., & Mortensen, E. (2003). Atlantic salmon *Salmo salar* L., brown trout *Salmo trutta* L. and Arctic charr *Salvelinus alpinus* (L.): a review of aspects of their life histories. *Ecology of Freshwater Fish*, 12(1), 1–59. doi:10.1034/j.1600-0633.2003.00010.x
- Knutsen, J. A., Knutsen, H., Gjosæter, J., & Jonsson, B. (2001). Food of anadromous brown trout at sea. *Journal of Fish Biology*, 59(3), 533–543. doi:10.1111/j.1095-8649.2001.tb02359.x
- Kristinsson, J. B. (1992). Aquaculture in Iceland—history and present status. *Icelandic Agricultural Science*, 6, 5–8.
- Krkošek, M., Lewis, M. A., & Volpe, J. P. (2005). Transmission dynamics of parasitic sea lice from farm to wild salmon. *Proceedings of the Royal Society B: Biological Sciences*, 272(1564), 689–696. doi:10.1098/rspb.2004.3027
- Krkošek, M., Morton, A., Volpe, J. P., & Lewis, M. A. (2009). Sea lice and salmon population dynamics: effects of exposure time for migratory fish. *Proceedings of the Royal Society B: Biological Sciences*, 276(1668), 2819–2828. doi:10.1098/rspb.2009.0317
- Krkošek, M. (2010). Host density thresholds and disease control for fisheries and aquaculture. *Aquaculture Environment Interactions*, 1(1), 21–32. doi:10.3354/aei0004
- Krkošek, M., Revie, C. W., Gargan, P. G., Skilbrei, O. T., Finstad, B., & Todd, C. D. (2013). Impact of parasites on salmon recruitment in the Northeast Atlantic Ocean. *Proceedings of the Royal Society B: Biological Sciences*, 280(1750), 20122359. doi:10.1098/rspb.2012.2359
- Kvenseth, P. G. (1997). Best current practice for lice control in Norway. *Caligus*, 2, 4–9.
- L'Abee-Lund, J. H., Jonsson, B., Jensen, A. J., Saettem, L. M., Heggberget, T. G., Johnsen, B. O., & Naesje, T. F. (1989). Latitudinal variation in life-history characteristics of sea-run migrant brown trout *Salmo trutta*. *The Journal of Animal Ecology*, 525–542.
- Largiader, C. R., Estoup, A., Lecerf, F., Champigneulle, A., & Guyomard, R. (2001). Microsatellite analysis of polyandry and spawning site competition in brown trout (*Salmo trutta* L.). *Genetics selection evolution*, 33, S205-S222.
- Lees, F., Gettinby, G., & Revie, C. W. (2008). Changes in epidemiological patterns of sea lice infestation on farmed Atlantic salmon, *Salmo salar* L., in Scotland between 1996 and 2006. *Journal of Fish Diseases*, 31(4), 259–268. doi:10.1111/j.1365-2761.2007.00897.x
- MacKinnon, B. 1997. Sea lice: a review. *World Aquaculture* 28:5–10.
- MacKinnon, B. (1998). Host factors important in sea lice infections. *ICES Journal of Marine Science*, 55(2), 188–192. doi:10.1006/jmsc.1997.0361
- MAST (2013). Declaration of disease-free status of compartments in accordance with Council Directive 2006/88/EC. Selfoss: MAST. Retrieved from <http://www.mast.is/english/library/Fr%C3%A6%C3%B0sluefni/DeclarationISAfreecompartments130323GJ.pdf>
- Matthews, M. A., Poole, W. R., Dillane, M. G., & Whelan, K. F. (1997). Juvenile recruitment and smolt output of brown trout (*Salmo trutta* L.) and Atlantic salmon (*Salmo salar* L.) from a lacustrine system in western Ireland. *Fisheries Research*, 31(1), 19–37. doi:10.1016/S0165-7836(97)00013-1
- McBeath, A. J., Penston, M. J., Snow, M., Cook, P. F., Bricknell, I. R., & Cunningham, C. O. (2006). Development and application of real-time PCR for specific detection of *Lepeophtheirus salmonis* and *Caligus elongatus* larvae in Scottish plankton samples. *Diseases of aquatic organisms*, 73(2), 141.
- McKenzie, E., Gettinby, G., McCart, K., & Revie, C. W. (2004). Time-series models of sea lice *Caligus elongatus* (Nordmann) abundance on Atlantic salmon *Salmo salar* L. in Loch Sunart, Scotland. *Aquaculture Research*, 35(8), 764–772. doi:10.1111/j.1365-2109.2004.01099.x
- McKibben, M. A., & Hay, D. W. (2004). Distributions of planktonic sea lice larvae *Lepeophtheirus salmonis* in the inter-tidal zone in Loch Torridon, Western Scotland in relation to salmon farm production cycles. *Aquaculture Research*, 35(8), 742–750. doi:10.1111/j.1365-2109.2004.01096.x
- McKinley, R. S., Finstad, B., Bjørn, P. A., & Hunter, K. J. (2003). *UBC Science Forum: Sea Lice Review and Summary of Research Priorities (Sea Lice White Paper)*.

- McLean, P. H., Smith, G. W., & Wilson, M. J. (1990). Residence time of the sea louse, *Lepeophtheirus salmonis* K., on Atlantic salmon, *Salmo salar* L., after immersion in fresh water. *Journal of Fish Biology*, 37(2), 311–314. doi:10.1111/j.1095-8649.1990.tb05861.x
- Middlemas, S. J., Fryer, R. J., Tulett, D., & Armstrong, J. D. (2013). Relationship between sea lice levels on sea trout and fish farm activity in western Scotland. *Fisheries Management and Ecology*, 20(1), 68–74. doi:10.1111/fme.12010
- Miller, S. H., & Morgan, S. G. (2013). Phenotypic plasticity in larval swimming behavior in estuarine and coastal crab populations. *Journal of Experimental Marine Biology and Ecology*, 449, 45–50. doi:10.1016/j.jembe.2013.08.013
- Mills, D. H. (1989). *Ecology and management of Atlantic salmon*. London, New York: Chapman and Hall.
- Moore, A., Lacroix, G. L., & Sturlaugsson, J. (2000). Tracking Atlantic salmon post-smolts in the sea. *The ocean life of atlantic salmon—environmental and biological factors influencing survival*, 49–64.
- Mordue A. J., & Birkett, M. A. (2009). A review of host finding behaviour in the parasitic sea louse, *Lepeophtheirus salmonis* (Caligidae: Copepoda). *Journal of Fish Diseases*, 32(1), 3–13. doi:10.1111/j.1365-2761.2008.01004.x
- Mustafa, A., Conboy, G. A., Burka, J. F., Hendry, C. I., & McGladdery, S. E. (2000). Life-span and reproductive capacity of sea lice, *Lepeophtheirus salmonis*, under laboratory conditions. *Special Publication-Aquaculture Association of Canada*, (4), 113–114.
- Mustafa, A., Rankaduwa, W., & Campbell, P. (2001). Estimating the cost of sea lice to salmon aquaculture in eastern Canada. *The Canadian Veterinary Journal*, 42(1), 54.
- NASCO (2011) Report of the twenty-eighth annual meeting of the council. North Atlantic Salmon Conservation Organization, Ilulissat, Greenland. pp. 1–349.
- Niemelä, E., Orell, P., Erkinaro, J., Dempson, J. B., Brørs, S., Svenning, M. A., & Hassinen, E. (2006). Previously spawned Atlantic salmon ascend a large subarctic river earlier than their maiden counterparts. *Journal of Fish Biology*, 69(4), 1151–1163. doi:10.1111/j.1095-8649.2006.01190.x
- O'Connell, M. F., & Dempson, J. B. (1996). Spatial and temporal distributions of salmonids in two ponds in Newfoundland, Canada. *Journal of Fish Biology*, 48(4), 738–757. doi:10.1111/j.1095-8649.1996.tb01469.x
- Oines, O., Simonsen, J. H., Knutsen, J. A., & Heuch, P. A. (2006). Host preference of adult *Caligus elongatus* Nordmann in the laboratory and its implications for Atlantic cod aquaculture. *Journal of Fish Diseases*, 29(3), 167–174. doi:10.1111/j.1365-2761.2006.00702.x
- O'Shea, B. A. (2005). Host identification and settlement of the infective copepodid stage of the salmon louse, *Lepeophtheirus salmonis* (Krøyer, 1837). *UNIVERSITY OF ABERDEEN (UNITED KINGDOM)*.
- Paisley, L. G., Ariel, E., Lyngstad, T., Jónsson, G., Vennerström, P., Hellström, A., & Østergaard, P. (2010). An overview of aquaculture in the nordic countries. *Journal of the World Aquaculture Society*, 41(1), 1–17. doi:10.1111/j.1749-7345.2009.00309.x
- Penston, M. J., & Davies, I. M. (2009). An assessment of salmon farms and wild salmonids as sources of *Lepeophtheirus salmonis* (Krøyer) copepodids in the water column in Loch Torridon, Scotland. *Journal of Fish Diseases*, 32(1), 75–88. doi:10.1111/j.1365-2761.2008.00986.x
- Penston, M. J., McKibben, M. A., Hay, D. W., & Gillibrand, P. A. (2004). Observations on open-water densities of sea lice larvae in Loch Shieldaig, Western Scotland. *Aquaculture Research*, 35(8), 793–805. doi:10.1111/j.1365-2109.2004.01102.x
- Penston, M. J., Millar, C. P., Zuur, A., & Davies, I. M. (2008). Spatial and temporal distribution of *Lepeophtheirus salmonis* (Krøyer) larvae in a sea loch containing Atlantic salmon, *Salmo salar* L., farms on the north-west coast of Scotland. *Journal of Fish Diseases*, 31(5), 361–371. doi:10.1111/j.1365-2761.2008.00915.x
- Pert, C. C., Fryer, R. J., Cook, P., Kilburn, R., McBeath, S., McBeath, A., . . . McCarthy, U. (2014). Using sentinel cages to estimate infestation pressure on salmonids from sea lice in Loch Shieldaig, Scotland. *Aquaculture Environment Interactions*, 5(1), 49–59.
- Piasecki, W. (1996). The developmental stages of *Caligus elongatus* von Nordmann, 1832 (Copepoda: Caligidae). *Canadian Journal of Zoology*, 74(8), 1459–1478.

- Pike, A. W., & Wadsworth, S. L. (2000). Sea lice on salmonids: Their biology and control. In *Advances in Parasitology. Advances in Parasitology Volume 44* (pp. 233–337). Elsevier.
- Price, P. W. (1980). Evolutionary biology of parasites: *Princeton University Press*.
- Rae, G. H. (1999). Sea lice, medicines and a national strategy for control. *Fish Veterinary Journal (United Kingdom)*,
- Rae, G. H. (2002). Sea louse control in Scotland, past and present. *Pest Management Science*, 58(6), 515–520. doi:10.1002/ps.491
- Revie, C. W., Dill, L., Finstad, B., & Todd, C. (2009). Sea lice working group report (Vol. 39). *Norsk institutt for naturforskning (NINA), Trondheim*.
- Revie, C. W., Gettinby, G., Treasurer, J. W., Rae, G. H., & Clark, N. (2002). Temporal, environmental and management factors influencing the epidemiological patterns of sea lice (*Lepeophtheirus salmonis*) infestations on farmed Atlantic salmon (*Salmo salar*) in Scotland. *Pest Management Science*, 58(6), 576–584. doi:10.1002/ps.476
- Revie, C. W., Gettinby, G., Treasurer, J. W., Wallace, C. (2003). Identifying epidemiological factors affecting sea lice *Lepeophtheirus salmonis* abundance on Scottish salmon farms using general linear models. *Diseases of aquatic organisms*, 57(1-2), 85.
- Rosten, T. W., Poulsen, H., Alanära, A., Eskelinen, U., Bergsson, A. B., & Olafsen, T. (2013). Perspectives for sustainable development of Nordic aquaculture: *Nordic Council of Ministers*.
- Salama, N. K. G., Collins, C. M., Fraser, J. G., Dunn, J., Pert, C. C., Murray, A. G., & Rabe, B. (2013). Development and assessment of a biophysical dispersal model for sea lice. *Journal of Fish Diseases*, 36(3), 323–337. doi:10.1111/jfd.12065
- Salama, N. K. G., & Murray, A. G. (2011). Farm size as a factor in hydrodynamic transmission of pathogens in aquaculture fish production. *Aquaculture Environment Interactions*, 2, 61–74.
- Schram, T. A. (2000). Practical identification of pelagic sea lice larvae. *Journal of the Marine Biological Association of the UK*, 84(1), 103–110. doi:10.1017/S0025315404008963h
- Schram, T. A. (2004). Practical identification of pelagic sea lice larvae. *Journal of the Marine Biological Association of the UK*, 84(01), 103-110.
- Serra-Llinares, R. M., Bjørn, P. A., Finstad, B., Nilsen, R., Harbitz, A., Berg, M., & Asplin, L. (2014). Salmon lice infection on wild salmonids in marine protected areas: an evaluation of the Norwegian ‘National Salmon Fjords’. *Aquaculture Environment Interactions*, 5(1), 1–16. doi:10.3354/aei00090
- Shearer, W. M. (1992). The Atlantic salmon: Natural history, exploitation and future management: *Fishing News Books*.
- Siegel, D. A., Kinlan, B. P., Gaylord, B., & Gaines, S. D. (2003). Lagrangian descriptions of marine larval dispersion. *Marine Ecology Progress Series*, 260, 83–96.
- Skilbrei, O. T., Finstad, B., Urdal, K., Bakke, G., Kroglund, F., & Strand, R. (2013). Impact of early salmon louse, *Lepeophtheirus salmonis* infestation and differences in survival and marine growth of sea-ranching Atlantic salmon, *Salmo salar* L., smolts 1997–2009. *Journal of Fish Diseases*, 36(3), 249–260. doi:10.1111/jfd.12052
- Stucchi, D. J., Guo, M., Foreman, M. G. G., Czajko, P., Galbraith, M., Mackas, D. L., & Gillibrand, P. A. (2011). Modelling sea lice production and concentrations in the Broughton Archipelago, British Columbia. *Salmon lice: An integrated approach to understanding parasite abundance and distribution*. Oxford, UK: Wiley-Blackwell,
- Sturlaugsson, J. & Johannsson, M. 1996. Migratory pattern of wild sea trout (*Salmo trutta* L.) in SE-Iceland recorded by data storage tags. *ICES C.M. 1996/M:5*
- Tapia, F. J., & Pineda, J. (2007). Stage-specific distribution of barnacle larvae in nearshore waters: potential for limited dispersal and high mortality rates. *Marine Ecology Progress Series*, 342, 177–190. doi:10.3354/meps342177
- Tchernavin, V. (1939). The origin of salmon: is its ancestry marine or freshwater. *Salmon and Trout Magazine*, 95, 120-40.

- Thorstad, E. B., Økland, F., Finstad, B., Sivertsgard, R., Bjørn, P. A., & McKinley, R. S. (2004). Migration speeds and orientation of Atlantic salmon and sea trout post-smolts in a Norwegian fjord system. *Environmental Biology of Fishes*, 71(3), 305–311. doi:10.1007/s10641-004-1264-7
- Thorstad, E. B., Økland, F., Finstad, B., Sivertsgård, R., Plantalech, N., Bjørn, P. A., & McKinley, R. S. (2007). Fjord migration and survival of wild and hatchery-reared Atlantic salmon and wild brown trout post-smolts. *Hydrobiologia*, 582(1), 99–107. doi:10.1007/s10750-006-0548-7
- Thorstad, E. B., Todd, C. D., Bjørn, P. A., Gargan, P. G., Vollset, K. W., Halattunen, E., . . . Finstad, B. (2014). Effects of salmon lice on sea trout - a literature review. *NINA Report* 1044, 1-162.
- Thorstad, E. B., Whoriskey, F., Uglem, I., Moore, A., Rikardsen, A. H., & Finstad, B. (2012). A critical life stage of the Atlantic salmon *Salmo salar*: behaviour and survival during the smolt and initial post-smolt migration. *Journal of Fish Biology*, 81(2), 500–542. doi:10.1111/j.1095-8649.2012.03370.x
- Tingley, G. (1997). The occurrence of lice on sea trout (*Salmo trutta* L.) captured in the sea off the East Anglian coast of England. *ICES Journal of Marine Science*, 54(6), 1120–1128. doi:10.1016/S1054-3139(97)80017-3
- Todd, C. D., Walker, A. M., Hoyle, J. E., Northcott, S. J., Walker, A. F., & Ritchie, M. G. (2000). Infestations of wild adult Atlantic salmon (*Salmo salar* L.) by the ectoparasitic copepod sea louse *Lepeophtheirus salmonis* Krøyer: prevalence, intensity and the spatial distribution of males and [2pt] females on the host fish. *Hydrobiologia*, 429(1/3), 181–196. doi:10.1023/A:1004031318505
- Torrissen, O., Jones, S., Asche, F., Guttormsen, A., Skilbrei, O. T., Nilsen, F., . . . Jackson, D. (2013). Salmon lice - impact on wild salmonids and salmon aquaculture. *Journal of Fish Diseases*, 36(3), 171–194. doi:10.1111/jfd.12061
- Treasurer, J. W. (2002). A review of potential pathogens of sea lice and the application of cleaner fish in biological control. *Pest management science*, 58(6), 546–558. doi:10.1002/ps.509
- Tucker, C. S., Sommerville, C., & Wootten, R. (2002). Does size really matter? Effects of fish surface area on the settlement and initial survival of *Lepeophtheirus salmonis*, an ectoparasite of Atlantic salmon *Salmo salar*. *Diseases of Aquatic Organisms*, 49(2), 145–152.
- Tully, O. (1989). The succession of generations and growth of the caligid copepods *Caligus elongatus* and *Lepeophtheirus salmonis* parasitising farmed Atlantic salmon smolts (*Salmo salar* L.). *Journal of the Marine Biological Association of the United Kingdom*, 69(02), 279. doi:10.1017/S0025315400029404
- Tully, O., Gargan, P., Poole, W. R., & Whelan, K. F. (1999). Spatial and temporal variation in the infestation of sea trout (*Salmo trutta* L.) by the caligid copepod *Lepeophtheirus salmonis* (Krøyer) in relation to sources of infection in Ireland. *Parasitology*, 119(01), 41–51. doi:10.1017/S003118209900445X
- Tully, O., & Nolan, D. T. (2002). A review of the population biology and host-parasite interactions of the sea louse *Lepeophtheirus salmonis* (Copepoda: Caligidae). *Parasitology*, 124(07). doi:10.1017/S0031182002001889
- Tully, O., & Whelan, K. F. (1993). Production of nauplii of *Lepeophtheirus salmonis* (Krøyer) (Copepoda: Caligidae) from farmed and wild salmon and its relation to the infestation of wild sea trout (*Salmo trutta* L.) off the west coast of Ireland in 1991. *Fisheries Research*, 17(1-2), 187–200. doi:10.1016/0165-7836(93)90018-3
- Urquhart, K., Pert, C. C., Kilburn, R., Fryer, R. J., & Bricknell, I. R. (2008). Prevalence, abundance, and distribution of *Lepeophtheirus salmonis* (Krøyer, 1837) and *Caligus elongatus* (Nordmann, 1832) on wild sea trout *Salmo trutta* L. *ICES Journal of Marine Science*, 65(2), 171–173. doi:10.1093/icesjms/fsm188
- Wagner, G. N., Fast, M. D., & Johnson, S. C. (2008). Physiology and immunology of *Lepeophtheirus salmonis* infections of salmonids. *Trends in Parasitology*, 24(4), 176–183. doi:10.1016/j.pt.2007.12.010
- Wagner, G. N., McKinley, R. S., Bjørn, P. A., & Finstad, B. (2003). Physiological impact of sea lice on swimming performance of Atlantic salmon. *Journal of Fish Biology*, 62(5), 1000–1009. doi:10.1046/j.1095-8649.2003.00091.x
- Wagner, G. N., McKinley, R. S., Bjørn, P. A., & Finstad, B. (2004). Short-term freshwater exposure benefits sea lice-infected Atlantic salmon. *Journal of Fish Biology*, 64(6), 1593–1604. doi:10.1111/j.0022-1112.2004.00414.x

- Ward, D. M., & Hvidsten, N. A. (2010). Predation: compensation and context dependence. *Atlantic salmon ecology*. Edited by Ø. Aas, A. Klemetsen, S. Einum, and J. Skurdal. Wiley-Blackwell, Oxford, UK, 199–220.
- Wells, A., Grierson, C. E., Marshall, L., MacKenzie, M., Russon, I. J., Reinardy, H., . . . Hazon, N. (2007). Physiological consequences of “premature freshwater return” for wild sea-run brown trout (*Salmo trutta*) postsmolts infested with sea lice (*Lepeophtheirus salmonis*). *Canadian Journal of Fisheries and Aquatic Sciences*, 64(10), 1360–1369. doi:10.1139/f07-107
- Wendelaar Bonga, S. E. (1997). The stress response in fish. *Physiological reviews*, 77(3), 591–625.
- Werner, E. E., & Gilliam, J. F. (1984). The ontogenetic niche and species interactions in size-structured populations. *Annual review of ecology and systematics*, 393–425.
- White, H. C. (1940). "Sea Lice" (*Lepeophtheirus*) and Death of Salmon. *Journal of the Fisheries Research Board of Canada*, 5a(2), 172–175. doi:10.1139/f40-017
- Wootten, R., Smith, J. W., & Needham, E. A. (1982). Aspects of the biology of the parasitic copepods *Lepeophtheirus salmonis* and *Caligus elongatus* on farmed salmonids, and their treatment. *Proceedings of the Royal Society of Edinburgh. Section B. Biological Sciences*, 81(03), 185–197. doi:10.1017/S0269727000003389