

# References

- Åberg, P. (1992). Size based demography of the seaweed *Ascophyllum nodosum* in stochastic environments. *Ecology*, 73: 1488-1501.
- Agriculture, Ministry of. (2017). Aquatic Plant Harvesting - Province of British Columbia [Government]. Retrieved May 22, 2017, from <http://www2.gov.bc.ca/gov/content/industry/agriculture-seafood/fisheries-and-aquaculture/commercial-fisheries/aquatic-plant-harvesting>
- Alban, F., Le Floc'h, P., & Boncoeur, J. (2004). The impact of economic and regulatory factors on the relative profitability of fishing boats: A case study of the seaweed harvesting fleet of Northwest Brittany (France). *Aquatic Living Resources*, 17(2), 185-193.
- Alexander, K. A., Potts, T. P., Freeman, S., Israel, D., Johansen, J., Kletou, D., ... & Angel, D. L. (2015). The implications of aquaculture policy and regulation for the development of integrated multi-trophic aquaculture in Europe. *Aquaculture*, 443, 16-23.
- Alþingi (2017). *Lög um breytingu á lögum um umgengni um nytjastofna sjávar, lögum um stjórn fiskveiða og lögum um veiðigjald (öflun sjávargróðurs í atvinnuskyni)*, Pub. L. No. 1048/146, 49 (2017).
- Andrade, A., Córdoba, R., Dave, R., Girot, P., Herrera-F., B., Munroe, R; Oglethorpe, J., Paaby, P., Pramova, E.; Watson, E.; Vergar, W., Suarez, I. (2012). Principles and Guidelines for Integrating Ecosystem-based Approaches to Adaptation in Project and Policy Design. *IUCN- CEM, CATIE*. Turrialba, Costa Rica. 4p.
- Ballesteros, E., Torras, X., Pinedo, S., García, M., Mangialajo, L., & de Torres, M. (2007). A new methodology based on littoral community cartography dominated by macroalgae for the implementation of the European Water Framework Directive. *Marine Pollution Bulletin*, 55(1-6), 172-180. <https://doi.org/10.1016/j.marpolbul.2006.08.038>
- Baardseth, E. M. (1955). Regrowth of *Ascophyllum nodosum* after harvesting. *Institute for Industrial Research and Standards*.
- Baardseth, E. M. (1970). Synopsis of biological data on knobbed wrack *Ascophyllum nodosum* (Linnaeus) Le Jolis. Rome: *Food and Agriculture Organization of the United Nations*.
- Barsh, R. L. (1997). Forests, indigenous people, and biodiversity. *Global Biodiversity*, 7:20-23.
- Benediktsson, K. (2014). Nature in the 'neoliberal laboratory.' *Dialogues in Human Geography*, 4(2), 141-146. <https://doi.org/10.1177/2043820614536340>

- Berkes, F., & Folke, C. (1998). *Linking social and ecological systems: Management practices and social mechanisms for building resilience*. Cambridge University Press, Cambridge, UK.
- Berkes, F., J. Colding, & C. Folke (eds.). (2003). *Navigating social-ecological systems: building resilience for complexity and change*. Cambridge University Press, Cambridge, UK.
- Bernard, H. R. (2006). *Research methods in anthropology: qualitative and quantitative approaches* / H. Russell Bernard (4th ed.). Lanham, Md: AltaMira Press.
- Bertness, M. D., & Leonard, G. H. (1997). The role of positive interactions in communities: lessons from intertidal habitats. *Ecology*, 78(7), 1976-1989.
- Bjarnason, M. T., Helgason, G. V., Eiríksson, T., & Jonsson, G. S. (2015). *Potential for harvesting and processing brown seaweed in Ísafjarðardjúp* (Business) (p. 18). Ísafjarðardjúp: Rorum.
- Black, R. & R.J. Miller. 1991. Use of the intertidal zone by fish in Nova Scotia. *Env. Biol. Fishes*, 31: 109–121.
- Boaden, P. J., & Dring, M. T. (1980). A quantitative evaluation of the effects of *Ascophyllum* harvesting on the littoral ecosystem. *Helgoländer Meeresuntersuchungen*, 33(1), 700.
- Breiðafjarðarnefnd. (2016, November 22). *Breiðafjarðarnefnd* [Environmental protection committee]. Retrieved February 15, 2018, from <http://www.breidafjordur.is/breidafjarðarnefnd/breidafjarðarnefnd/>
- Buschmann, A. H., Camus, C., Infante, J., Neori, A., Israel, Á., Hernández-González, M. C., ... & Critchley, A. T. (2017). Seaweed production: overview of the global state of exploitation, farming and emerging research activity. *European Journal of Phycology*, 52(4), 391-406.
- [CAFF] Conservation of Arctic Flora and Fauna. (2002). *Protected Areas of the Arctic: Conserving a Full Range of Values*. Ottawa, 2002. pp.16
- [CBD] Convention on Biological Diversity (2007). In-depth review of the application of the ecosystem approach. *Barriers to the application of the ecosystem approach*. In: *Proceedings of the 12th meeting of the subsidiary body on scientific, technical and technological advice*. Paris: UNESCO
- Cottier-Cook, E. J., Nagabhatla, N., Badis, Y., Campbell, M. L., Chopin, T., Dai, W., ... & Huo, Y. (2016). Safeguarding the future of the global seaweed aquaculture industry. *United Nations University (INWEH) and Scottish Association for Marine Science Policy Brief*. p. 12.
- Cousens, R. (1982). Popular misconceptions about *Ascophyllum nodosum* and their effect on the interpretation of its population dynamics from existing data. *Br. Phycol. J.*, 17: 231.

- Christie, H., Jørgensen, N., Norderhaug, K., & Waage-Nielsen, E. (2003). Species distribution and habitat exploitation of fauna associated with kelp (*Laminaria Hyperborea*) along the Norwegian Coast. Marine Biological Association of the United Kingdom. *Journal of the Marine Biological Association of the United Kingdom*, 83(4), 687–699.
- Chu, S. H., Zhang, Q. S., Liu, S. K., Tang, Y. Z., Zhang, S. B., Lu, Z. C., & Yu, Y. Q. (2012). Tolerance of *Sargassum thunbergii* germlings to thermal, osmotic and desiccation stress. *Aquatic botany*, 96(1), 1-6.
- Davison, I. R. & Pearson, G. A. (1996). Stress tolerance in intertidal seaweeds. *J. Phycol.* 32:197–211.
- Dayton, P., Currie, V., Gerrodette, T., Keller, B., Rosenthal, R., Tresca, D. (1984) Patch dynamics and stability of California kelp communities. *Ecol Monogr* 54:253–289
- DFO. (1999). The impact of the rockweed harvest on the habitat of southwest New Brunswick. *DFO Maritimes Regional Habitat Status Report*, 99/2E.
- Douve, F. (2008). The importance of marine spatial planning in advancing ecosystem-based sea use management. *Marine Policy*, 32(5), 762–771. Retrieved from: <https://doi.org/https://doi.org/10.1016/j.marpol.2008.03.021>
- Dudgeon, S., & Petraitis, P. S. (2005). First year demography of the foundation species, *Ascophyllum nodosum*, and its community implications. *Oikos*, 109(2), 405–415. <https://doi.org/10.1111/j.0030-1299.2005.13782.x>
- Durrenberger, E. P., & Pálsson, G. (1987). Ownership at Sea: Fishing Territories and Access to Sea Resources. *American Ethnologist*, 14(3), 508–522.
- Eckersley, L. K., & Scrosati, R. A. (2012). Temperature, desiccation, and species performance trends along an intertidal elevation gradient. *Curr Dev Oceanogr*, 5(5), 59-73.
- Edwards, A. (1980). Ecological studies of the kelp, *Laminaria hyperborea*, and its associated fauna in south-west Ireland. *Ophelia*, 19(1), 47-60.
- FAO. (1987). *Case Studies of Seven Commercial Seaweed Resources (Technical Paper No. 281)* (p. 311). Rome. Retrieved from <http://www.fao.org/docrep/x5819e/x5819e00.htm>
- FAO (2010). *Resources, technologies and services for future aquaculture: a needs assessment for sustainable development*. Retrieved from <http://agris.fao.org/agris-search/search.do?recordID=AV2012045157>
- FAO. (2014). *Seaweeds and other aquatic plants: Capture Production 2014 (FAO Yearbook of Fishery and Aquaculture Statistics)*. FAO.
- FAO. (2016). *The State of World Fisheries and Aquaculture 2016. (Contributing to food security and nutrition for all.)* (p. 200). Rome: FAO.

- Fegley, J. C. (2001). Ecological implications of rockweed, *Ascophyllum Nodosum* (L.) le jolis, harvesting. *Electronic Theses and Dissertations*. 397. <http://digitalcommons.library.umaine.edu/etd/397>
- Ferguson, M. A., Williamson, R. G., & Messier, F. (1998). Inuit knowledge of long-term changes in a population of Arctic tundra caribou. *Arctic*, 201-219.
- Fleurence, J. (1999). Seaweed proteins: biochemical, nutritional aspects and potential uses. *Trends in Food Science & Technology*, 10(1), 25–28. [https://doi.org/10.1016/S0924-2244\(99\)00015-1](https://doi.org/10.1016/S0924-2244(99)00015-1)
- FMC. (2018). FMC Corporation - Corporate Overview [Corporation]. Retrieved January 16, 2018, from <http://www.fmc.com/About-FMC/Corporate-Overview>
- Gilchrist, G., Mallory, M., & Merkel, F. (2005). Can local ecological knowledge contribute to wildlife management? Case studies of migratory birds. *Ecology and Society*, 10(1), n/a. <https://doi.org/10.5751/ES-01275-100120>
- Graham, M. H., Vásquez, J. A., Buschmann A. H. (2007). Global ecology of the giant kelp *Macrocystis*: from ecotypes to ecosystems. *Ocean Mar Biol Ann Rev* 45:39–88
- Government Offices of Iceland. (2018, February 14). Organizational Chart of the Ministry of Industries and Innovation [Government]. Retrieved February 15, 2018, from <https://www.government.is/ministries/ministry-of-industries-and-innovation/organizational-chart/>
- Guiry, M. (2017, May). Seaweed.ie :: Information on marine algae [Biology]. Retrieved November 21, 2016, from <http://www.seaweed.ie/aquaculture/WorldProduction.php>
- Gunn, A., Arlooktoo, G., and Kaomayok, D. (1988). The contribution of ecological knowledge of Inuit to wildlife management in the Northwest Territories. Pages 22-30 in M. M. R. Freeman and L. N. Carbyn, editors. *Traditional knowledge and renewable resource management in northern regions*. IUCN Commission on Ecology and the Boreal Institute for Northern Studies, Occasional Publication Number 23, Edmonton, Alberta, Canada.
- Gunnarsson, K., Gunnar, J., & Pálsson, Ó. (1998). *Sjávarnyttjar við Ísland*. Reykjavík: Mál og menning.
- Gunnarsson, K., Bourgos, J., Gunnarsdóttir, L., Egilsdóttir, S., Georgsdóttir, G. I., & Madrigal, V. F. P. (2017). *Klóbang í Breiðafirði útbreiðsla og magn*. Iceland: Hafrannsóknastofnun.
- Gunnarsson, K., Jónsson, G., & Pálsson, Ó. K. (1998). *Sjávarnyttjar við Ísland. Mál og menning*.
- Harley, C. D., Anderson, K. M., Demes, K. W., Jorve, J. P., Kordas, R. L., Coyle, T. A., & Graham, M. H. (2012). Effects of climate change on global seaweed communities. *Journal of Phycology*, 48(5), 1064-1078.

- Hoey, D. (2017, March 21). Judge rules that rockweed harvesters need landowners' permission. *Portland Press Herald*. Retrieved May 26, 2017, from <http://www.pressherald.com/2017/03/21/judge-rules-that-rockweed-harvesters-need-to-obtain-landowners-permission/>
- Icelandic Institute of Natural History. (2018, March 4). Klóþangsfjörur [Scientific research institute]. Retrieved March 4, 2018, from <https://www.ni.is/greinar/klothangsfjorur>
- Icelandic Road and Coastal Administration. (2017, August 22). Tides [Weather and sea state information]. Retrieved February 23, 2018, from <https://tinyurl.com/y8cr6w3r>
- Jenkins, S.R., Norton T.A. & Hawkins, S.J. (1999) Direct and indirect effects of a macroalgal canopy and limpet grazing in structuring a sheltered intertidal community. *Marine Ecology Progress Series 188*: 81–92.
- Johannes, R.E. (1998). The case for data-less marine resource management: examples from tropical nearshore fisheries. *Trends in Ecology and Evolution 13*:243-246.
- Johnson, B. (1999). The Role of Adaptive Management as an Operational Approach for Resource Management Agencies. *Conservation Ecology*, 3(2). <https://doi.org/10.5751/ES-00136-030208>
- Johnson, S.C., Scheibling, R.E. (1987). Structure and dynamics of epifaunal assemblages on intertidal macroalgae *Ascophyllum nodosum* and *Fucus vesiculosus* in Nova Scotia, Canada. *Mar. Ecol. Prog. Ser.* 37, 209-227.
- Jonsson, G. S., Helgason, G. V., Eiríksson, T., Bjarnason, M. T., Agustsson, T., & Provan, F. (2015). *Feasibility of harvesting brown seaweed from the Ísafjardardjup area*. Rorum.
- Juanich, G. L. (1988). *Manual on seaweed farming 1. Eucheuma spp. (Technical Paper No. ASEAN/SF/88/Manual No. 2)*. Manila, Philippines: ASEAN/UNDP/FAO Regional Small-Scale Coastal Fisheries Development Project. Retrieved from <http://www.fao.org/docrep/field/003/AC416E/AC416E00.htm>
- Kelly, L., Collier, L., Costello, M. J., Diver, M., McGarvey, S., Kraan, S., ... & Giury, M. D. (2001). *Impact assessment of hand and mechanical harvesting of Ascophyllum nodosum on regeneration and biodiversity*.
- Keser, M. & Larson B.R. (1984). Colonization and growth of *Ascophyllum nodosum* (Phaeophyta) in Maine. *J. Phycol.*, 20, pp. 83-87.
- Kiely Cove Marine Enterprises. (2016). A Review of Historical Harvests and Utilization of *Ascophyllum nodosum* in Breiðafjörður Bay in Support of Sustainable Management, Draft Report to Thorverk Reykhólar Iceland (Draft report) (p. 42). Iceland: Kiely Cove Marine Enterprises.
- Lazo, L., & Chapman, A. R. O. (1996). Effects of harvesting on *Ascophyllum nodosum* (L.) Le Jol. (Fucales, Phaeophyta): a demographic approach. *Journal of Applied Phycology*, 8(2), 87–103. <https://doi.org/10.1007/BF02186311>

- Lee, B. (2008). Seaweed - Potential as a marine vegetable and other opportunities. Rural Industries Research and Development Corporation (RIRDC) Publication no. 08/009, Kingston, ACT, Australia. [rirdc.infoservices.com.au/downloads/08-009](http://rirdc.infoservices.com.au/downloads/08-009) p.34
- Leonard, G. H. (2000). Latitudinal variation in species interactions: a test in the New England rocky intertidal zone. *Ecology*, 81(4), 1015-1030.
- Leslie, H. M., & McLeod, K. L. (2007). Confronting the challenges of implementing marine ecosystem-based management. *Frontiers in Ecology and the Environment*, 5(10), 540-548.
- Ludwig, D. (1993). Environmental sustainability: magic, science, and religion in natural resource management. *Ecological Applications*, 3(4), 555-558.
- Lubchenco, J. (1980). Algal zonation in the New England rocky intertidal community: an experimental analysis. *Ecology*, 61(2), 333-344.
- Maack, M. (2018). Personal communication March 2, 2018. Project manager, Vestfjarðastofu, Reykhólar
- Maack, M. (2018). Personal communication March 18, 2018. Project manager, Vestfjarðastofu, Reykhólar
- Mackinson, S., & Nottestad, L. (1998). Points of view: combining local and scientific knowledge. *Reviews in Fish Biology and Fisheries*, 8(4), 481-490.
- Maine Department of Marine Resources (2017). Commercial landings. Retrieved February 1, 2017, from <https://www.maine.gov/dmr/commercial-fishing/landings/documents/12-16LandingsBySpeciesWithBonus.Table.pdf>
- MarBEF. (2013, August 16). European Marine Biodiversity Research Sites [Marine biodiversity data system]. Retrieved January 23, 2018, from <http://www.marbef.org/data/sitedetails.php?id=12933>
- Mathieson, A. C., Penniman, C. A., Busse, P. K., and E. Tveter-Gallagher (1982). Effects of ice on *Ascophyllum nodosum* within the Great Bay estuary system of New Hampshire Maine. *J. Phycol.* 18: 331-336.
- McGinnis, M., & Ostrom, E. (2014). Social-ecological system framework: initial changes and continuing challenges. *Ecology and Society*, 19(2). <https://doi.org/10.5751/ES-06387-190230>
- McHugh, D. (2003). A Guide to the Seaweed Industry. FAO Technical Paper 441. Food and Agricultural Organisation of the United Nations, Rome. [www.fao.org/docrep/006/y4765e/y4765e00.htm](http://www.fao.org/docrep/006/y4765e/y4765e00.htm)
- McLeod, K.L., Lubchenco, J., Palumbi S., Rosenberg A. (2005). Scientific Consensus Statement on Marine Ecosystem-Based Management. *Communication Partnership for Science and the Sea*.

- Miura, A. (1975). Porphyra cultivation in Japan. In: Tokida and Hirose. *Advance of Phycology in Japan*. VEB G. Fischer Verlag, Jena, Germany. pp. 273-304.
- Monagle, M. M., Cornish, M. L., Morrison, L., Araújo, R., & Critchley, A. (2017). Sustainable harvesting of wild seaweed resources. *European Journal of Phycology*, 52(4), 371–390. <https://doi.org/DOI: 10.1080/09670262.2017.1365273>
- Mouritsen, O. G., Dawczynski, C., Duelund, L., Jahreis, G., Vetter, W., & Schröder, M. (2013). On the human consumption of the red seaweed dulse (*Palmaria palmata* (L.) Weber & Mohr). *Journal of Applied Phycology*, 25(6), 1777-1791.
- National Research Council. (2002). *The drama of the commons*. National Academies Press.
- Nayar, S., & Bott, K. (2014). Current status of global cultivated seaweed production and markets. *World Aquaculture*. Retrieved from [https://www.researchgate.net/publication/265518689\\_Current\\_status\\_of\\_global\\_cultivated\\_seaweed\\_production\\_and\\_markets](https://www.researchgate.net/publication/265518689_Current_status_of_global_cultivated_seaweed_production_and_markets)
- Naylor, J. (1976). Production, trade and utilization of seaweeds and seaweed products. *FAO Fisheries Technical Papers (FAO)*. Documents Techniques FAO sur les Peches (FAO)-Documentos Tecnicos de la FAO sobre la Pesca (FAO). no. 159.
- NOAA. (2017). NOAA's National Ocean Service: Tide cycle variations [Government]. Retrieved December 20, 2017, from [https://oceanservice.noaa.gov/education/kits/tides/media/supp\\_tide07a.html](https://oceanservice.noaa.gov/education/kits/tides/media/supp_tide07a.html)
- National Oceanic and Atmospheric Administration [NOAA]. (2017). What is seaweed? [Government]. Retrieved May 22, 2017, from <http://oceanservice.noaa.gov/facts/seaweed.html>
- NOAA Fisheries. (2007, June 18). Local Fisheries Knowledge Project [Government]. Retrieved July 28, 2017, from [https://www.st.nmfs.noaa.gov/lfkproject/02\\_c.definitions.htm](https://www.st.nmfs.noaa.gov/lfkproject/02_c.definitions.htm)
- Nunes, N., Ferraz, S., Valente, S., Barreto, M. C., & Carvalho, M. A. A. P. de. (2017). Biochemical composition, nutritional value, and antioxidant properties of seven seaweed species from the Madeira Archipelago. *Journal of Applied Phycology*, 29(5), 2427–2437. <https://doi.org/10.1007/s10811-017-1074-x>
- Ólafsdóttir, S. (2009, September). Icelandic Coastal Zone Management in Global Perspective. Coastal Zone Management presented at the Symposium on Coastal Zone Management and Spatial Marine Planning University Centre of the Westfjords, Iceland. Retrieved from [https://www.uw.is/haskolasetur\\_vestfjarda/skraarsafn/skra/268/](https://www.uw.is/haskolasetur_vestfjarda/skraarsafn/skra/268/)
- Olsen, J. L., Zechman, F. W., Hoarau, G., Coyer, J. A., Stam, W. T., Valero, M., & Åberg, P. (2010). The phylogeographic architecture of the furoid seaweed *Ascophyllum nodosum*: an intertidal 'marine tree' and survivor of more than one glacial–interglacial cycle. *Journal of Biogeography*, 37(5), 842-856.

- Ostrom, E. (2009). A General Framework for Analyzing Sustainability of Social-Ecological Systems. *Science*, 325(5939), 419–422. <https://doi.org/10.1126/science.1172133>
- Ostrom, E., Gardner, R., & Walker, J. (1994). *Rules, games, and common-pool resources*. University of Michigan Press.
- Pálmason, S. G. (Producer), Magnason A. S. & Thorfinnur, G. (Directors). (2009). *Dreamland* [Motion picture]. Iceland: Ground Control Productions.
- Pálsson, G., & Durrenberger, E. P. (1996). *Images of Contemporary Iceland: Everyday Lives and Global Contexts*. University of Iowa Press. Retrieved from <https://muse.jhu.edu/book/12500>
- Peckol, P., Harlin, M. M., & Krumscheid, P. (1988). Physiological and Population Ecology of Intertidal and Subtidal *Ascophyllum Nodosum* (Phaeophyta) 1. *Journal of Phycology*, 24(2), 192-198.
- Petraitis, P. S., & Latham, R. E. (1999). The importance of scale in testing the origins of alternative community states. *Ecology*, 80(2), 429-442
- Petit, R. J., & Hampe, A. (2006). Some evolutionary consequences of being a tree. *Annu. Rev. Ecol. Evol. Syst.*, 37, 187-214.
- Phillippi, A., Tran, K., & Perna, A. (2014). Does intertidal canopy removal of *Ascophyllum nodosum* alter the community structure beneath?. *Journal of experimental marine biology and ecology*, 461, 53-60.
- POC. Pew Oceans Commission 2003. *America's Living Oceans: Charting a Course for Sea Change*. Arlington (VA).
- Pollnac, R. B., Abbott-Jamieson, S., Smith, C., Miller, M. L., Clay, P. M., & Oles, B. (2006). Toward a model for fisheries social impact assessment. *Marine Fisheries Review*, 68(1-4), 1-18.
- Rainer, R., (1997). A Plan for Community Management of the St. Croix Estuary Area. Rockweed Harvesting. St. Andrews, New Brunswick.
- Raymond, C. M., Fazey, I., Reed, M. S., Stringer, L. C., Robinson, G. M., & Evely, A. C. (2010). Integrating local and scientific knowledge for environmental management. *Journal of Environmental Management*, 91(8), 1766–1777. <https://doi.org/10.1016/j.jenvman.2010.03.023>
- Rebours, C., Marinho-Soriano, E., Zertuche-González, J. A., Hayashi, L., Vásquez, J. A., Kradolfer, P., ... & Robledo, D. (2014). Seaweeds: an opportunity for wealth and sustainable livelihood for coastal communities. *Journal of Applied Phycology*, 26(5), 1939–1951. <https://doi.org/10.1007/s10811-014-0304-8>
- Rebours, C., & Meland, M. (2012). Introduction to the management and regulation of the Norwegian seaweed industry. *Bioforsk FOKUS*, 7(2), 278–279.



- Rebours, C., & Meland, M. (2012). The Norwegian Seaweed Industry Work Package 1 & 2. Netalgae. Retrieved from [http://www.netalgae.eu/uploadedfiles/Norwegian\\_seaweed\\_industry\\_WP12.pdf](http://www.netalgae.eu/uploadedfiles/Norwegian_seaweed_industry_WP12.pdf)
- Reykjólahreppur. (2018, March 4). Reykjólahreppur [Government]. Retrieved March 4, 2018, from [http://www.reykholar.is/stjornsysla/skrifstofa\\_reykholahrepps/](http://www.reykholar.is/stjornsysla/skrifstofa_reykholahrepps/)
- Reykjólaskóli. (2018, March 4). Reykjólaskóli [Education Institution]. Retrieved March 4, 2018, from <http://www.reykholar.is/skoli/skolinn/starfsfolk/>
- Rist, L., Felton, A., Samuelsson, L., Sandström, C., & Rosvall, O. (2013). A New Paradigm for Adaptive Management. *Ecology and Society*, 18, 63. <https://doi.org/10.5751/ES-06183-180463>
- Rothman, M. D., Anderson, R. J., & Smit, A. J. (2006). The effects of harvesting of the South African kelp (*Ecklonia maxima*) on kelp population structure, growth rate and recruitment. In *Eighteenth International Seaweed Symposium* (pp. 109-115). Springer, Dordrecht.
- Samband Íslenskra Sveitarfélaga. (2017, January 1). Reykjólahreppur [Government Statistics]. Retrieved February 9, 2018, from <http://www.samband.is/sveitarfelogin/vefdir/reykholahreppur/>
- Sarà, G., Pirro, M. D., Romano, C., Rumolo, P., Sprovieri, M., & Mazzola, A. (2007). Sources of organic matter for intertidal consumers on *Ascophyllum*-shores (SW Iceland): a multi-stable isotope approach. *Helgoland Marine Research*, 61(4), 297–302. <https://doi.org/10.1007/s10152-007-0078-3>
- Schmidt, A. L., Coll, M., Romanuk, T. N., & Lotze, H. K. (2011). Ecosystem structure and services in eelgrass *Zostera marina* and rockweed *Ascophyllum nodosum* habitats. *Marine Ecology Progress Series*, 437, 51-68.
- Schonbeck, M. W., & Norton, T. A. (1980). Factors controlling the lower limits of furoid algae on the shore. *Journal of Experimental Marine Biology and Ecology*, 43(2), 131-150.
- Scottish Government, Marine Scotland, & ABP Marine Environmental Research. (2016). *Wild Seaweed Harvesting Strategic Environmental Assessment Report (Consultation Paper)* (p. 263). Scotland: Scottish Government. Retrieved from <https://beta.gov.scot/publications/wild-seaweed-harvesting-strategic-environmental-assessment-environmental-report/>
- SeaCell. (2017). Smartfiber AG: SeaCell LT. Retrieved May 22, 2017, from <http://www.smartfiber.de/en/fibers/seacelltm-lt/>
- Seeley, R. H., & Schlesinger, W. H. (2012). Sustainable seaweed cutting? The rockweed (*Ascophyllum nodosum*) industry of Maine and the Maritime Provinces. *Annals of the New York Academy of Sciences*, 1249(1), 84-103.

- Sharp, G. J., & Pringle, J. D. (1990). Ecological impact of marine plant harvesting in the northwest Atlantic: a review. *Hydrobiologia*, 204(1), 17-24.
- Sharp, G. (1987). *Ascophyllum nodosum and its harvesting in Eastern Canada (Technical Paper)*. Rome: FAO. Retrieved from <http://www.fao.org/docrep/x5819e/x5819e04.htm>
- Sjómælingar Íslands, 2016. *Sjávarfallatöflur 2016*. Sjómælingadeild Landhelgisgæslunnar, Reykjavík. 27p.
- Sjötun, K. (1993). Seasonal lamina growth in two age groups of *Laminaria saccharina* (L.) Lamour. in western Norway. *Botanica marina*, 36(5), 433-442.
- Stagnol, D., Renaud, M., & Davoult, D. (2013). Effects of commercial harvesting of intertidal macroalgae on ecosystem biodiversity and functioning. *Estuarine, Coastal and Shelf Science*, 130, 99-110.
- Statistics Iceland. (2017). Iceland. Retrieved from [http://px.hagstofa.is/pxen/pxweb/en/Ibuar/Ibuar\\_\\_mannfjoldi\\_\\_1\\_yfirlit\\_\\_yfirlit\\_mannfjolda/MAN00000.px/table/tableViewLayout1/?rxid=691568a8-dc14-40d5-ab2b-8dee3ae6f258](http://px.hagstofa.is/pxen/pxweb/en/Ibuar/Ibuar__mannfjoldi__1_yfirlit__yfirlit_mannfjolda/MAN00000.px/table/tableViewLayout1/?rxid=691568a8-dc14-40d5-ab2b-8dee3ae6f258)
- Stengel, D., & Dring, M. (1997). Morphology and in situ growth rates of plants of *Ascophyllum nodosum* (Phaeophyta) from different shore levels and responses of plants to vertical transplantation. *European Journal of Phycology*, 32(2), 193–202. <https://doi.org/10.1080/09670269710001737129>
- Sharp, G. J., Ugarte, R., & Semple, R. (2006). The ecological impact of marine plant harvesting in the Canadian Maritimes, Implications for coastal zone management. *Science Asia*, 32(2006), 77-86. doi:10.2306/scienceasia1513-1874.2006.32(s1).077
- Stelma, B. (2015). Adding the fourth dimension to 3D printing. ExtremeTech.com; New York.
- Stévant, P., Rebours, C., & Chapman, A. (2017). Seaweed aquaculture in Norway: recent industrial developments and future perspectives. *Aquaculture International*, 1–18. <https://doi.org/10.1007/s10499-017-0120-7>
- Thorverk HF. (2018, January 3). Thorverk policy; Excellence based on safety [Company]. Retrieved February 13, 2018, from [https://www.thorverk.is/about\\_us/the\\_company/](https://www.thorverk.is/about_us/the_company/)
- Ugarte, R. (2007). Review of the management of rockweed (*Ascophyllum nodosum*) harvesting in New Brunswick after a decade of its initiation. In: Pohle G. W., Wells P. G., & Rolston S. J. (eds.) Challenges in Environmental Management in the Bay of Fundy-Gulf of Maine. Proceedings of the 7th Bay of Fundy Science Workshop, St. Andrews, New Brunswick, 24–27 October 2006. *Bay of Fundy Ecosystem Partnership Technical Report No. 3*. Bay of Fundy Ecosystem Partnership, Wolfville, NS. N° 3: 108-116

- Ugarte, R. (2011). An evaluation of the mortality of the brown seaweed *Ascophyllum nodosum* (L.) Le Jol. produced by cutter rake harvests in southern New Brunswick, Canada. *J. Appl. Phycol.* 23: 401-407.
- Ugarte, R., & Sharp, G. (2012). Management and production of the brown algae *Ascophyllum nodosum* in the Canadian maritimes. *Journal of applied phycology*, 24(3), 409-416.
- UN General Assembly. (2015, October 21). *Transforming our world: the 2030 Agenda for Sustainable Development*. UN General Assembly. Retrieved from [http://www.un.org/ga/search/view\\_doc.asp?symbol=A/RES/70/1&Lang=E](http://www.un.org/ga/search/view_doc.asp?symbol=A/RES/70/1&Lang=E)
- [USCOP] US Commission on Ocean Policy. (2004). *An Ocean Blueprint for the 21st Century*. USCOP.
- Vadas, R. L., Wright, W. A., & Beal, B. F. (2004). Biomass and Productivity of Intertidal Rockweeds (*Ascophyllum nodosum* LeJolis) in Cobscook Bay. *Northeastern Naturalist*, 11, 123–142.
- Vandermeulen, H. (2013). *Information to Support Assessment of Stock Status of Commercially Harvested Species of Marine Plants in Nova Scotia: Irish Moss, Rockweed and Kelp*. Department of Fisheries and Oceans, Maritimes Region. Retrieved from [http://publications.gc.ca/collections/collection\\_2013/mpo-dfo/Fs70-5-2013-042-eng.pdf](http://publications.gc.ca/collections/collection_2013/mpo-dfo/Fs70-5-2013-042-eng.pdf)
- Vásquez, J. A., Piaget, N., & Vega, J. M. A. (2012). The *Lessonia nigrescens* fishery in northern Chile: “how you harvest is more important than how much you harvest.” *Journal of Applied Phycology*, 24(3), 417–426. <https://doi.org/10.1007/s10811-012-9794-4>
- Walters, C. J., & Hilborn, R. (1978). Ecological optimization and adaptive management. *Annual review of Ecology and Systematics*, 9(1), 157-188.
- WCED. (1987). *Report of the World Commission on Environment and Development: Our Common Future*. UN-WCED. Retrieved from <http://www.un-documents.net/wced-ocf.htm>
- West-Iceland Institute of Natural History. (2004, June). *Breiðafjörður Nature and History*. The Breiðafjörður Committee
- West-Iceland Institute of Natural History. (2006, April). *Birdlife in Breiðafjörður*. The Breiðafjörður Committee
- WFD. (2000). Directive 2000/60/EC of the European Parliament and of the Council establishing a framework for the Community action in the field of water policy. *Official Journal of the European Communities*. Retrieved from <http://eur-lex.europa.eu/legal-content/EN/TXT/?uri=CELEX:32000L0060>. Accessed: January 16, 2018.

- White, H., & Glenn, H. (2006). Environmental impact mitigation and bi-culture: a comparative legal analysis of flexibility within European legal regimes – biofilter deployment. *Aquaculture International*, 14(3), 297–317. <https://doi.org/10.1007/s10499-005-9035-9>
- Witman, J. D., Etter, R. J., & Smith, F. (2004). The relationship between regional and local species diversity in marine benthic communities: A global perspective. *Proceedings of the National Academy of Sciences*, 101(44), 15664–15669. <https://doi.org/10.1073/pnas.0404300101>
- Zabel, C. J., Roberts, L. M., Mulder, B. S., Stauffer, H. B., Dunk, J. R., Wolcott, K., ... & Goldsmith, G. (2002). A collaborative approach in adaptive management at a large landscape scale. *Predicting species occurrence: issues of accuracy and scale*. Predicting species occurrence: issues of accuracy and scale, 241-253. Island Press, Washington, D.C., USA.
- Zemke-White, W. L., & Ohno, M. (1999). World seaweed utilisation: An end-of-century summary. *Journal of Applied Phycology*, 11(4), 369–376. <https://doi.org/10.1023/A:1008197610793>
- 1107/145 *Stjórnarfrumvarp: umgengni um nytjastofna sjávar o.fl.*, Pub. L. No. 1107 (2015). Retrieved from <http://www.althingi.is/altext/145/s/1107.html>